

VOL I
FORMIC ACID AND DERIVATIVES #97

Volume I Formic Acid & Derivatives

#97

1/9/74

VOLUME 1

GRAS MONOGRAPH SERIES FORMIC ACID AND DERIVATIVES

**prepared for
THE FOOD AND DRUG ADMINISTRATION
DEPARTMENT OF HEALTH, EDUCATION
AND WELFARE**

JANUARY 9, 1974

This publication was prepared under Contract Number FDA 72-100
with the Public Health Service, Food and Drug Administration,
Department of Health, Education, and Welfare

**prepared by
Tracor Jitco, Inc.**

Table of Contents

SUMMARY

CHEMICAL INFORMATION

	<u>Page</u>
I. Nomenclature	1
II. Empirical Formula	1
III. Structural Formula	1
IV. Molecular Weight	1
V. Specifications	1
VI. Description	2
VII. Analytical Methods	3
VIII. Occurrence and Levels	7

BIOLOGICAL DATA

I. Acute Toxicity	9
A. Fish	9
B. Daphnias	9
C. Mice	9
D. Rats	9
E. Guinea Pigs	13
F. Cats	13
G. Dogs	13
H. Human Toxicity	14
II. Short-term Studies	15
A. Yeasts	15
B. Rats	17
C. Guinea Pigs	23
D. Rabbits	23
E. Dogs	27
F. Sheep	27
G. Cattle	28
H. Human Toxicity	32
III. Long-term Studies	35
A. Rats	35
IV. Special Studies	37
A. Bacteria	37
B. Insects	39
C. Chick Embryos	39
D. Mice	42

BIOCHEMICAL ASPECTS

	<u>Page</u>
I. Breakdown	43
II. Absorption-Distribution	43
III. Metabolism and Excretion	43
A. Oxidation	43
B. Excretion	45
C. Intermediary Metabolism	48
IV. Effects on Enzymes and Other Biochemical Parameters	50
V. Drug Interaction	52
VI. Consumer Exposure Information	53

BIBLIOGRAPHY

JOURNAL ARTICLE COPIES & TRANSLATIONS

SUMMARY

Description

Formic acid is a colorless, highly corrosive, fuming liquid with a sharp, penetrating odor. It is very soluble in water, ethanol, ether, and glycerol.

Ethyl formate is a colorless, unstable liquid with a pleasant peach-like odor and a slightly bitter taste. It is miscible with ethanol, ether, and benzene. It is also slightly soluble in water, but decomposes. Ethyl formate is highly flammable and a dangerous fire and explosion risk.

Sodium formate is a white, deliquescent solid with a slight formic acid odor. It is soluble in water and glycerol, slightly soluble in ethanol, but insoluble in ether.

Occurrence

Formic acid occurs naturally in the poison of ants, wasps, bees, and some other insects. It is a normal constituent of mammalian muscle tissue, sweat, and urine. A number of plants such as nettles, unripe grapes, peaches, raspberries, etc. also contain formic acid. Relatively large amounts are formed when coffee is roasted. Formic acid is present also in wines, rum, and certain other distilled fermented fruit beverages. It is found in some mineral waters. Reported natural concentrations range from 2 to 770 mg per 100 grams.

Ethyl formate is found naturally in a variety of plant oils, fruits, and juices such as oranges, apples, pears, and in honey. It is present also in wines, rum, and certain other distilled liquors. Apparently it does not occur naturally in the animal kingdom. In the food industry, ethyl formate is used as a flavoring in candy,

chewing gum, ice cream, baked goods, etc., and in many essences. It is used also as a fumigant for certain dried fruits, nuts, cereals, and tobacco. (018,139,491). Possible daily intake values range from 1 to 84 mg approximately.

Formic acid and sodium formate are included in the category of substances possibly migrating to foods from paper and paperboard food containers, where they are used in manufacturing processes.

Acute Toxicity

Formic acid in mice has an oral LD₅₀ values of 1100 mg/kg BW (275), 940 mg/kg via the peritoneum (435), and 145 mg/kg when injected intravenously (275); when administered intrarectally, the lethal dose is 1250 mg/kg BW (491). In rats, the oral LD₅₀ is 1830 mg/kg BW (435); in rabbits the oral MLD is greater than 4000 mg/kg (491) while the intravenous MLD is 239 mg/kg (433). For daphnias and fish, the LD₅₀ values are 70 and 175 mg/liter, respectively (109,435).

Ethyl formate has an oral LD₅₀ of 1850 mg/kg BW in rats (217) and a value of 1110 mg/kg in guinea pigs (217). By inhalation, the fatal dose is 24 mg/liter in rats (429) and 32 mg/liter in cats (491).

Sodium formate has an oral LD₅₀ of 11,200 mg/kg BW in mice (275) and an MLD of 4000 mg/kg in dogs (420). In mice, by the intravenous route, the LD₅₀ is 807 mg/kg (275); in dogs the MLD is 3000 mg/kg (430). For fish, the LD₅₀ is 5000 mg/liter by immersion (109).

Short-term studies

Formic acid is reported to have caused anemia, leucopenia, basophilic neutropenia, and a slight lymphocytosis in rats when given at a level of 1% in the drinking water. The offspring mortality rate also was stated to be high (425). In another study, formic acid at a level of 360 mg/kg BW daily for 9 weeks, after 90 mg/kg BW daily for

17 weeks, depressed food intake and retarded growth (430). Weanling rats fed diets containing ethyl formate as high as 10,000 ppm for 17 weeks grew normally and remained in good health throughout. Hematologic findings were normal and no gross or microscopic abnormalities were detected (173).

Guinea pigs and rabbits exposed to ethyl formate in the atmosphere at levels of 40 to 130 mg per liter suffered marked central nervous system depression (260). On the other hand, doses as large as 1000 mg/kg BW subcutaneously were tolerated without any evidence of toxicity (260,491). Silage containing formic acid at a level of 1/2 gallon per ton, fed to cattle for 12 weeks, apparently caused no harmful effects (074).

In man, massive doses of sodium formate (2-3 grams several times a day) caused vertigo, nausea, vomiting, albuminuria, hematuria, vesical tenesmus, dyspnea, and lowered temperature (445). A single dose of 3-4 grams caused albuminuria and hematuria in one of three subjects but the condition cleared up after five days (491). Daily doses of 150 mg/kg BW per os for some time caused no harm (430). A single dose of 4 grams intravenously was tolerated without signs of toxicity (491). Formic acid in 500 mg amounts in lemonade given daily for 4 weeks caused no harmful effects (491). Larger doses produced irritation of the gastrointestinal tract followed by vomiting and diarrhea (491).

Formic acid and formates are quite toxic via the respiratory tract. Inhalation of concentrations as low as 32 mg per liter by human subjects caused progressive irritation of the eye and mucous membranes which persisted for as long as four hours (491). Locally, formic acid is corrosive to the skin and mucous membranes. Even mild exposures may be followed by severe after-effects (491).

Long-term Studies

Rats tolerated sodium formate at a level of 730 mg/kg BW daily in the drinking water for 1.5 years without evidence of toxic effects (275).

Special Studies

Formic acid at a level of 0.1% caused mutation in Drosophila melanogaster (450). In concentrations as low as 0.005%, it induced mutation in Escherichia coli from streptomycin-dependence to non-dependence (097).

Doses of sodium formates as high as 20 mg per egg were not toxic or teratogenic for chick embryos (275).

No statistically significant evidence of histologic abnormalities were detected in the skin of mice painted with 8% formic acid for a period of 50 days (142).

Biochemistry

Formic acid and sodium formate are rapidly absorbed from the stomach. Formic acid is absorbed also through the intact skin and via the lungs. Ethyl formate is absorbed through the lungs as well as from the gastrointestinal tract (271,491).

Absorbed formic acid or formate apparently is partly oxidized, partly excreted unchanged in the urine, and partly metabolized in the tissues. The liver appears to be the main site of formate oxidation (491). The oxidation mechanism, in the rat at least, involves a catalase-hydrogen peroxide complex and the enzymes responsible for peroxide formation (331). Folic acid and vitamin B₁₂ are essential for normal formate oxidation (197,320). Small doses of formic acid administered to experimental animals are usually oxidized almost completely whereas 50% or more of larger doses are excreted unchanged in the urine (491). The biological half-life varies according to the species, from 12 minutes in the rat to 77 minutes in dogs. Values of 45-46 minutes have been obtained with human subjects (275,277).

Formic acid is a normal constituent of human urine and individuals on a mixed diet excrete from 13 to 120 mg daily (156,277). The fate of ethyl formate in the body apparently has not been determined (491).

The participation of the formate ion in intermediary metabolism is well-established (011,156). Retained formate is rapidly utilized in the formation of protein, lipids, carbohydrates, and nucleic acids and these in turn are distributed throughout all tissues of the body (370,434). Formate is a precursor, through one-carbon metabolism, of serine, methionine, cysteine, and purines; it ultimately becomes incorporated through these precursors into proteins and nucleic acids (370). Formate is incorporated also into carbohydrates via glucose and the glucogenic amino acid serine (011).

Normally, only small amounts of formate are ingested in foods but many compounds contribute to the C_1 pool for formation of endogenous formate; among these are serine, glycine, histidine, methionine, choline, sarcosine, and acetone (056).

Folic acid, vitamin E, and ascorbic acid are involved in control of formate metabolism; deficiencies in any of these cause metabolic derangement (105,197,320).

The nervous system appears to be especially susceptible to the action of formates. Small doses cause depression; large amounts cause convulsions, paralysis of the medulla, and death (491).

Heart action is stimulated by trace amounts (1 ppm) of formic acid. Concentrations as low as 1:10,000, however, quickly and irreversibly stop heart action. Symptoms of cardiac injury are increase and then decrease of the systole, slowing of the heart beat, contraction irregularities, arrhythmias, and diastole arrest (491).

Formic acid intravenously causes immediate vasoconstriction. Sodium formate, on the other hand, causes vasodilation in the liver, brain, and kidney, but vasoconstriction in the limbs (491).

Gastric secretion and peristalsis are stimulated by formic acid. There is also a diuretic effect. Single large doses or repeated small doses cause kidney damage (491).

Formic acid inhibits lysozyme, ribonuclease, and trypsin (425). Small doses cause a persistent methemoglobinuria apparently due to action on catalase.

Formic acid is reported to have strong germicidal properties (491) and has been used as a food preservative in certain European countries (137). Germicidal action is believed to reside in the un-ionized molecule (491). Methyl, ethyl, and isopropyl formates were effective against various life cycle stages of the dried fruit beetle, raisin moth, and other insects that infest dried fruits, nuts, cereals, and tobacco. They are used as fumigants in these areas of the food industry (018,139,485,491).

Folic acid antagonist drugs such as methotrexate inhibit metabolism of formic acid, resulting in increased excretion of it in the urine (275).

Consumer Exposure

Formic acid is found in a number of plants, fruits, fruit juices, and wines as well as in milk, meats, certain vegetables, and some mineral waters (056,146,275).

In the food industry, formic acid is used to a limited extent as a flavor adjunct in candy, ices, ice cream, baked goods and several other food categories (146).

Ethyl formate occurs naturally in certain fruits, fruit juices, wines, rum, and some other distilled alcoholic beverages (146). It is a component of a number of food flavorings and is used in chewing gum, candy, ice cream,

ices, baked goods, and several other food categories (139). Maximum daily intakes up to 84 mg have been reported (138). Ethyl formate is used also as a bulk and package fumigant for certain dried fruits, nuts, cereals, and tobacco (018,139,491). With raisins and dried Zante currants, the maximum concentration permitted in the final product is 250 ppm (018).

Formic acid and sodium formate are included in the category of substances possibly migrating to foods from paper and paperboard food containers that employ formates in their manufacture (019).

Safe levels

Acceptable daily intake levels for formic acid and ethyl formate are 0-5 mg/kg body weight, for each, calculated as total formic acid and from all food additive sources. Maximum allowable concentrations per 8-hour work shift are 9 mg per cubic meter of air for ethyl formate (016,515,516).

CHEMICAL INFORMATION

I. Nomenclature

A. Common name:

Formic acid: Formic acid; Methanoic acid; Hydrogen carboxylic acid

Ethyl formate: Ethyl formate; Formic ether; Ethyl methanoate

Sodium formate: Sodium formate

B. Chemical name

Formic acid: Formic acid

Ethyl formate: Formic acid, ethyl ester

Sodium formate: Formic acid, sodium salt

C. Trade names:

Formic acid: Formisoton; Acirufan; Formidium (Therapeutic preparations)

D. Chemical Abstracts Services Unique Registry Number:

Formic acid: 64186

Ethyl formate: 109944

Sodium formate: 141537

II. Empirical formula

Formic acid: HCO_2H

Ethyl formate: $\text{HCO}_2\text{C}_2\text{H}_5$

Sodium formate: NaCHO_2

III. Structural formula

Formic acid: $\begin{array}{c} \text{O} \\ // \\ \text{H}-\text{C}-\text{OH} \end{array}$

Ethyl formate: $\begin{array}{c} \text{O} \quad \text{H} \quad \text{H} \\ // \quad | \quad | \\ \text{H}-\text{C}-\text{O}-\text{C}-\text{C}-\text{H} \\ \quad | \quad | \\ \quad \text{H} \quad \text{H} \end{array}$

Sodium formate: $\begin{array}{c} \text{O} \\ // \\ \text{H}-\text{C}-\text{O}-\text{Na} \end{array}$

IV. Molecular Weight

Formic acid: 46.03

Ethyl formate: 74.08

Sodium formate: 68.01

V. Specifications

The Food Chemicals Codex (088) lists the following specifications for food-grade formic acid and ethyl formate.

Formic acid

Assay - not less than 85 percent of CH_2O_2

Dilution test - passes test.

Limits of impurities

Acetic acid - not more than 0.4 percent

Arsenic (as AS) - not more than 3 ppm (0.0003%)

Heavy metals (as Pb) - not more than 10 ppm (0.001%)

Sulfate - not more than 40 ppm (0.004%)

Ethyl formate

Assay - not less than 95 percent $\text{C}_3\text{H}_6\text{O}_2$

Refractive index - between 1.359 and 1.363 at 20°C.

Solubility in alcohol - passes test

Specific gravity - between 0.916 and 0.921

Limits of impurity

Free acid (as formic acid) - not more than 0.1%

Arsenic (as As) - not more than 3 ppm (0.0003%)

Heavy metals (as Pb) - not more than 10 ppm (0.001%)

Sulfate - not more than 40 ppm (0.004%)

VI. Description

A. Formic acid - a colorless, highly corrosive, fuming liquid with a pungent, penetrating odor.

Physical constants:

Solubility: Completely miscible with water, ethanol, ether, and glycerol.

Specific gravity: 1.2201

Refractive index: 1.3719

Melting point: 8.3°C

Boiling point: 100.8°C

Flash point: 156°F.

Autoignition temperature: 1114°F.

May deteriorate in normal storage causing a hazard.

B. Ethyl formate - a colorless, unstable liquid with a pleasant, peach-like odor and a slightly bitter taste.

Physical constants:

Solubility: Miscible with ethanol, ether, and benzene.
Slightly soluble in water but decomposes.
Refractive index: 1.35975
Melting point: -80.5°C.
Boiling point: -54.3°C.
Flash point: -4°F.
Autoignition temperature: 851°F.
Is highly flammable and a dangerous fire and explosion risk.

C. Sodium formate - a white, deliquescent solid with a slight formic acid odor.

Physical constants:

Solubility: Soluble in water (1:13 RT) and glycerol;
slightly soluble in ethanol; insoluble in
ether.
Specific gravity: 1.92
Melting point: 253°C.
Is deliquescent; containers must be tightly closed.

VII. Analytical Methods

A. The Association of Official Agricultural Chemists (AOAC) has presented a steam distillation method for the extraction of volatile fatty acids from a sample (027). The apparatus used is shown in Figure 1. Any similar distillation assembly can be used if it is designed to handle the same volumes as the AOAC assembly, and $57 \pm 2\%$ recovery of acetic acid is obtained upon distillation.




Figure 1. Steam Distillation Assembly of the AOAC (027)

B. Formic acid may be detected by formation of derivatives. The melting points of some characteristic derivatives are given in Table 1. Similar derivatives of other volatile fatty acids are included for comparison (080).

Table 1. Derivatives of Some Volatile Fatty Acids and Their Melting Points (080).

Acid	B.P. (°C)	M.P. (°C)	p-toluidide	Derivatives and melting points		p-nitro- benzyl ester
				anilide	p-bromo- phenacyl ester	
Formic	100.7	8.4	53	50	140;135	31
Acetic	118.2	16.6	153;147	114	86.0	78
Propionic	141	-20.8	126;123	106	63.4	31
n-Butyric	162.5; 164	-5.5; -8	75	96;97	63	35
n-Pentanoic	186.4	-34.5	74	63	75	---

C. When formic acid is present as the only source of acidity in the sample, or when other acids are present in negligible amounts, a standard alkaline titration will suffice to measure the formic acid content. Titration with 0.05 N NaOH to a phenolphthalein end-point is a standard method (015).

D. Where other acids which do not react readily with permanganate are present, a redox titration may be used for formic acid determination (015). The formic acid solution is made alkaline with sodium carbonate and permanganate is added. The solution is acidified, an excess of oxalic acid is added, and a back titration performed. The redox titration may also be performed by adding excess permanganate to the alkaline sample and heating to 70°C. A sufficient amount of KI solution is added, followed by a back-titration using standard $\text{Na}_2\text{S}_2\text{O}_3$ (thiosulfate) (015).

E. Formic acid may be determined in the presence of other acids (notably acetic acid) by its reduction of soluble mercury(II)chloride, HgCl_2 to insoluble mercury(I)chloride, Hg_2Cl_2 . The reaction requires two hours of heating under a reflux condenser. The mercury is filtered and dried, and the weight used to calculate formic acid content of the original sample (015). This method is also presented by the AOAC for formic acid determination following steam distillation of the sample (027,028).

F. Shelley, et al. (420) have presented a method for the quantitative measurement of formic acid and other volatile fatty acids by gas chromatography. The method employs methyl enanthoate as an internal standard. Recoveries ranged from 95-105% for lower concentrations of acids and 97-101% for higher concentrations. The method was compared to the "Official" AOAC method (reduction of mercury(II)chloride, above) and good agreement was found. The gas chromatographic method was found to be more sensitive to the lower concentrations of acids. AOAC methods (028) included a gas chromatographic method for formic acid and the other volatile fatty acids similar to the method of Shelley, et al.

Watson and Crescuolo (603) found that microgram and sub-microgram quantities of formic, acetic, and propionic acids may be determined by gas chromatography as the p-substituted benzyl esters. Optimum results were realized with p-bromobenzyl derivatives. Table 2 shows the retention time of the various p-substituted benzyl esters; Fig. 2 is a chromatograph of a micro preparation of the p-bromobenzyl esters.

Table 2. Retention Times (min) of p-substituted Benzyl Formate Acetate and Propionate Esters and Other Relevant Related Compounds (503)

	p-Methylbenzyl (column temp. 100°)	p-Bromobenzyl (column temp. 120°)	p-Nitrobenzyl (column temp. 145°)
Formate	8.4	10.9	11.0
Acetate	15.1	17.0	15.3
Propionate	26.3	27.4	22.8
Alcohol	6.6	10.2	12.6
Halide	9.3	12.7	25.5
Artifact	5.7	7.5	7.2




Figure 2. Chromatograph of p-Bromobenzyl Ester Reaction Mixture (micro preparation using three-component acid solution (503)).

VIII. Occurrence and Levels

A. Formic acid

Formic acid occurs in the poison of ants (Formicidae), bees, wasps, stinging caterpillars, etc. and is partly responsible for its irritant effects. It was discovered in 1670 by distilling red ants (Formica rufa) with water. Bees impart some of their formic acid to the honey. Certain of the Isopoda contain formic acid and have diuretic properties (445). Formic acid is also a normal constituent of mammalian muscle tissue, sweat, and urine (275).

A number of plants contain formic acid. Nettles, Douglas fir needles, unripe grapes, petitgrain lemons, peaches, raspberries, strawberries, mace oil, valerian root extract, and bitter orange are typical examples (139,146). Relatively large amounts are formed when coffee is roasted and this is concentrated in coffee extracts (146). Formic acid is present also in rum, wines, and certain other distilled fermented fruit beverages. It is found in some mineral waters (156).

Formic acid is used to a certain extent industrially. It is a rubber latex coagulant, a decalcifier, reducer in dyeing and finishing of textiles and paper, and a constituent of certain sizes (438). It is used in dehairing and plumping hides, tanning, reclaiming old rubber, ore flotation, and electroplating (146). In the food industry, it is used as a flavoring adjunct, animal feed additive, brewing antiseptic, and, in certain European countries, as a food preservative (137,146,156). Levels found in some common foods and beverages are shown in Table 3.

Table 3. Formic Acid Content Reported for Some Common Foods and Beverages

Food	Amount Formic Acid (mg/100g)	Reference
Fruits	2-4	414
Fruit juices	3-10	501
Fruit syrups	65-163	406
Honey	2-200	128
Wines	0.1-34	464,475
Coffee, roasted	135-220	098
Coffee extracts	200-770	098
Milk (evap.)	3-4	295
Cheese	2-30	210,255

B. Ethyl formate

Ethyl formate is found in a variety of plant fruits and oils. Oranges, apples, pears, honey, and the oil of Boronia dentigeroides, are typical examples. It is present also in wines and certain distilled liquors such as rum (146).

No reports were found of its presence in the animal kingdom.

Ethyl formate is used industrially in the manufacture of cellulose acetate and nitrate and as a solvent for oils, fats, and waxes. In the food industry, it is used as a flavoring agent and as a fumigant (fungicide and larvacide for certain dried fruits, nuts, cereals, and tobacco (018,139,491)).

C. Sodium formate

No information was found concerning the occurrence of sodium formate in plants or animals. Commercially, it is used in adhesives, as a reducing agent, as a mordant, in the dyeing and printing of fabrics, and in tanning. Chemically, it is used as a buffer and as a solubilizer for trivalent metallic ions.

BIOLOGICAL DATA

I. Acute Toxicity

A. Fish

Dowden et al (109) exposed Bluegills (number, age, sex, weight not given) to graded amounts of formic acid or sodium formate in an organic-free medium for a period of 24 hours and determined the median tolerance (TL_m), i.e., the minimum concentration lethal to 50% of the fish within the 24 hour exposure period. The results are given in Table 4.

B. Daphnias

Sporn et al (435) subjected Daphnias to varying levels of formic acid in a nutrient medium over a period of 48 hours. (See Table 4.)

C. Mice

1. Sporn et al (435) injected white mice (number, age, weight, sex, and strain not given) intraperitoneally with varying amounts of formic acid in an acute toxicity study. (See Table 4.)

2. Malorny (275) determined the oral and intravenous LD₅₀ values for formic acid and several of its salts (see Tables 4 and 5) with white mice (age, weight, sex, and strain not specified), employing 45 to 55 animals for each compound.

D. Rats

1. Jenner, Fitzhugh et al (217) determined the oral (by intubation) LD₅₀ value for ethyl formate in young adult Osborne-Mendel rats, 10 per group evenly divided by sex, after a fasting period of 18 hours. Water was accessible at all times. Food was returned after administration of the compound. The animals were closely observed for toxic symptoms and the time of death recorded for all fatalities. Observations were continued until the survivors appeared normal and showed weight gain. The data are represented in Tables 4 and 6.

2. Sporn et al (435) determined the oral LD₅₀ (see table 4) of formic acid in young white rats (number, weight, sex, and strain not specified). Dyspnea and cyanosis preceded death. Weakness and diarrhea with a slow return to normal was characteristic of the survivors.

3. Smyth et al (429) exposed rats, six per group, (strain, age, sex, and weight not given), to ethyl formate vapor of varying concentrations for a period of four hours and recorded the fatalities over a 14-day observation period. The results are summarized in Table 4.

BIOLOGICAL DATA

I. Acute Toxicity

A. Fish

Dowden et al (109) exposed Bluegills (number, age, sex, weight not given) to graded amounts of formic acid or sodium formate in an organic-free medium for a period of 24 hours and determined the median tolerance (TLM), i.e., the minimum concentration lethal to 50% of the fish within the 24 hour exposure period. The results are given in Table 4.

B. Daphnias

Sporn et al (435) subjected Daphnias to varying levels of formic acid in a nutrient medium over a period of 48 hours. (See Table 4.)

C. Mice

1. Sporn et al (435) injected white mice (number, age, weight, sex, and strain not given) intraperitoneally with varying amounts of formic acid in an acute toxicity study. (See Table 4.)

2. Malorny (275) determined the oral and intravenous LD₅₀ values for formic acid and several of its salts (see Tables 4 and 5) with white mice (age, weight, sex, and strain not specified), employing 45 to 55 animals for each compound.

D. Rats

1. Jenner, Fitzhugh et al. (217) determined the oral (by intubation) LD₅₀ value for ethyl formate in young adult Osborne-Mendel rats, 10 per group evenly divided by sex, after a fasting period of 18 hours. Water was accessible at all times. Food was returned after administration of the compound. The animals were closely observed for toxic symptoms and the time of death recorded for all fatalities. Observations were continued until the survivors appeared normal and showed weight gain. The data are represented in Tables 4 and 6.

2. Sporn et al. (435) determined the oral LD₅₀ (see table 4) of formic acid in young white rats (number, weight, sex, and strain not specified). Dyspnea and cyanosis preceded death. Weakness and diarrhea with a slow return to normal was characteristic of the survivors.

3. Smyth et al. (429) exposed rats, six per group, (strain, age, sex, and weight not given), to ethyl formate vapor of varying concentrations for a period of four hours and recorded the fatalities over a 14-day observation period. The results are summarized in Table 4.

Table 4. Acute Toxicity of Formic Acid, Ethyl Formate, and Sodium Formate

Substance	Animal	Sex & No.	Route	Dosage mg/kg	Measurement	Reference/ Bibliography No.
Formic Acid	Fish	-	Immersion	175 ^a	LD ₅₀	Dowden (109)
Formic Acid	Daphnias	-	Immersion	70 ^a	LD ₅₀	Sporn (435)
Formic Acid	Mice	-	p.o.	1100	LD ₅₀	Malorny (275)
Formic Acid	Mice	-	i.v.	145	LD ₅₀	Malorny (275)
Formic Acid	Mice	-	i.p.	940	LD ₅₀	Sporn (435)
Formic Acid	Mice	-	i.r.	1250	LD	von Oettingen (491)
Formic Acid	Rats	-	p.o.	1830	LD ₅₀	Sporn (435)
Formic Acid	Rabbits	-	p.o.	4000+	MLD	von Oettingen (491)
Formic Acid	Rabbits	-	i.v.	239	MLD	Spector (433)
Ethyl Formate	Rats	-	p.o.	1850	LD ₅₀	Jenner (217)
Ethyl Formate	Rats	-	Inhalation	24 ^a	FD	Smyth (429)
Ethyl Formate	Guinea pigs	-	p.o.	1110	LD ₅₀	Jenner (217)
Ethyl Formate	Cats	-	Inhalation	32 ^a	LD	von Oettingen (491)

^a mg/liter

Table 4. (Continued)

Substance	Animal	Sex & No.	Route	Dosage mg/kg	Measurement	Reference Bibliography No.	
Sodium Formate	Fish	-	Immersion	5000 ^a	LD ₅₀	Dowden	(109)
Sodium Formate	Mice	-	p.o.	11,200	LD ₅₀	Malorny	(275)
Sodium Formate	Mice	-	i.v.	807	LD ₅₀	Malorny	(279)
Sodium Formate	Dogs	-	p.o.	4000	MLD	Sollmann	(430)
Sodium Formate	Dogs	-	i.v.	3000	MLD	Sollmann	(430)

12 ^a mg/liter

Table 5. LD₅₀ Formic Acid and its Salts (Mouse) (275)

Substance	LD ₅₀ mg/kg	LD ₅₀ Range mg/kg	Calculated according to formate	Number of animals
Per os				
Formic acid	1100	1000-1200	1076	55
Sodium formate	11200	9600-12800	7410	45
Potassium formate	5500	5000-6000	2950	50
Ammonium formate	2250	2050-2460		
Calcium formate	1920	1280-2560	1330	45
Intravenous				
Formic acid	145	138-151	142	50
Sodium formate	807	800-813	534	50
Potassium formate	95	93-97	51	55
Ammonium formate	410	408-412	293	50
Calcium formate	154	150-158	107	55

Table 6. Acute Oral Toxicity of Ethyl Formate (217)

Animal	LD ₅₀ (mg/kg)	Slope function	Toxic signs and death times (D.T)
Rat	1850	1.6	Depression within 5-10 minutes.
	(1520-2240) ^a	(1.3-1.9) ^a	Labored respiration. D.T. 15 min-2 hr.
Guinea pig	1110 ^a	2.0	Depression. Irritated
	(887-1390)	(1.3-3.1) ^a	gastrointestinal tract. D.T. 10 min-2 hr.

^a 95% confidence limits

E. Guinea Pigs

1. Jenner, Fitzhugh et al. (217) administered varying doses of ethyl formate to groups of young guinea pigs consisting of males and females, by intubation, after a fasting period of 18 hours. The animals were then given free access to food and water, and closely observed for signs of toxicity. The time of death was recorded for all fatalities and observation continued until survivors appeared normal and gained weight. The LD₅₀ value, slope function and their confidence limits along with toxic symptoms and death times are given in Tables 4 and 6.

2. von Oettingen (491) reported that according to Duquenois and Revel, inhalation by guinea pigs of air saturated with ethyl formate vapors "resulted within a few minutes in tremors, progressive depression of the central nervous system, and death from circulatory and respiratory failure."

F. Cats

Flury and Neuman (260) exposed cats (number, strain, age, weight and sex not specified) to several concentrations of ethyl formate vapor in air for varying periods of time. The animals were then closely observed for toxic signs and the time of death recorded for all fatalities (see Table 4).

Concentrations as low as 32 mg/liter for 20 minutes caused moderate irritation of the mucous membranes of the eye and respiratory tract. Progressive narcosis resulted after 80 minutes, and death with pulmonary edema followed an exposure period of 90 minutes.

A concentration of 44 mg/liter caused marked irritation of the mucous membranes, severe dyspnea, and staggering after 17 minutes. When the exposure time was increased to 20 minutes and then discontinued, one of the two animals recovered but the other died from pulmonary edema.

G. Dogs

1. von Oettingen (491), reporting an early study by Fleig, states that the fatal dose of 4000 mg/kg BW of formic acid for the dog per os causes repeated vomiting, clonic convulsions, progressive dyspnea prior to death by respiratory paralysis.

2. Stern (445) reports that formic acid diminishes heart action and blood pressure. The fatal dose was 1000 mg/kg BW (route not specified). Death was due to cardiac paralysis.

H. Human Toxicity

Reviews and surveys indicate that severe and fatal cases of formic acid poisoning in man are not unusual (491). Thirty-four cases were reported over an eight-year period in Travancore, the largest rubber-producing area in India, where the acid is extensively used for the coagulation of latex. Twenty-seven fatalities were suicidal, six accidental, and one due to an over-dose. Although the minimum amount ingested in these cases was not available, two ounces of the concentrated acid were always fatal. Quantities detected in the victims ranged from 0.04 to 31.8 grams (anhydrous acid) (222).

The signs and symptoms of fatal formic acid poisoning in man are: salivation, burning sensation in the mouth and pharynx, agonizing pain, bloody vomitus, and diarrhea. The pulse and respiration are rapid and soft at first but then slow down. The skin becomes cold and clammy, blood pressure drops and shock ensues. There is respiratory distress, cyanosis, and death may follow within 48 hours. Some patients develop albuminuria (with casts), hematuria, subsequent anuria, and may die from uremia, circulatory failure, or pneumonia (491).

Pathologic changes after ingestion of formic acid are similar to those of other corrosive acids. The tongue, palate, pharynx, and esophagus become hyperemic, swollen, and covered with necrotic spots. The larynx and trachea may be edematous and inflamed with hemorrhages. Stomach walls may be edematous with necrotic areas and contain bloody material or they may be dry and brittle. The jejunum is usually like the stomach but less severe. The kidneys are hyperemic and may be hemorrhaged and resemble the state in nephritis. The liver may show hemosiderin deposits (491).

The minimum fatal dose of formic acid and ethyl formate for man by mouth is estimated to be 30 grams or approximately 500 mg/kg BW (433). Sodium formate is less toxic with estimates ranging from 1000 to 4000 mg/kg BW orally. (155,430). The formate ion, however, may have a direct action on the brain in sublethal concentrations (491). Methemoglobinemia also has been detected in experimental animals (491). Formic acid and ethyl formate are toxic via the respiratory tract also and maximum allowable concentrations for eight hours of exposure daily are nine and 300 mg per cubic meter of air, respectively (016).

II. Short-term Studies

A. Yeasts

Watts (504) studied the action of formic acid, methanol, and formaldehyde on the oxidation of glucose, succinate, and ascorbate by yeast and rat liver slices and homogenates in an effort to shed light on the mechanism of methanol poisoning. Conventional Warburg manometric techniques were used. Results are shown in Fig. 3 and Tables 7 and 8.

Various enzymes were inhibited (3-10%) by formic acid at a level as low as 0.005M (Table 8). Tissue respiration was depressed 50% by concentrations as low as 0.04 to 0.16M (Table 7 and Fig. 3). These concentrations approximate the level found in tissues during fatal methanol poisoning in the rabbit with the minimum lethal dose. Further study showed, however, that formaldehyde (another tissue oxidation product of methanol) was ten times as potent as formic acid as a respiration inhibitor. The author concluded, therefore, that the former was more likely the main toxic factor in methanol poisoning rather than formic acid which had been postulated.




Fig. 3 The Inhibition of Oxidation of Succinate by Rat Liver Homogenate by Increasing Concentrations of Methanol, Formaldehyde, and Formic Acid (504).

Table 7. Concentration of Methanol, Formaldehyde, and Formic Acid Required to Produce 50% Inhibition of Tissue Respiration (504)

Tissue	Substrate and activity measured	Methanol M	Concentration Formaldehyde M	Formic acid M
Yeast	Glucose oxidation	>1.5	0.003	0.32
Yeast	Glucose fermentation	>1.0	0.003	0.16
Liver slices	Glucose oxidation	>1.0	0.010	0.50
Liver homogenate	Succinate oxidation	3.2	0.003	0.04
Liver homogenate	Ascorbate oxidation	>1.5	0.010	0.07

Table 8. The Comparative Inhibition of Tissue Respiration by 0.005 M Methanol, Formaldehyde and Formic Acid (504)

Tissue ^a and determination	Substrate	Control mm. ³ gas. mean \pm s.e.	Methanol %	Change from control Formaldehyde %	Formic acid %
Yeast, Q_{O_2} , (wet weight)	Glucose	79.0 \pm 2.4	+0.09	-56.6 ^b	+10.1
Yeast, Q_{O_2} (wet weight)	Glucose	29.3 \pm 0.5	-9.9	-67.3 ^b	-5.1 ^c
Liver slices, Q_{O_2} (wet weight)	Glucose	1.47 \pm 0.07	-0.7	-39.8 ^b	-8.2
Liver homogenate Q_{O_2} (wet weight)	Succinate	23.6 \pm 0.5	-6.6	-60.5 ^b	-10.3 ^c
Liver homogenate Q_{O_2} (wet weight)	Ascorbate	15.7 \pm 1.4	-1.3	-24.8 ^b	-3.2

^a Additions to Warburg vessels:

(a) Five milligrams of yeast was suspended in 2 ml. of 0.02 M KH_2PO_4 (pH 4.8) to which was added 1 ml. of 3% glucose. Temperature 30°C.

(b) Liver slices were suspended in Krebs-Ringer phosphate containing 0.3% glucose. Temperature 37°C.

(c) With minor modifications the homogenate method as described by Potter (5) was used for the oxidation of succinate and ascorbate.

(d) The final volume in all vessels was 3.0 ml. plus 0.2 ml. 20% KOH in center well for oxygen uptake measurements. Formic acid was adjusted to the pH of the suspension media before adding it to the vessels.

^b Statistical analysis of the data gave *P* values of less than 0.001 for these changes; this indicates the inhibition is highly significant.

^c *P* values for these changes were less than 0.05 which indicates they are probably significant changes. *P* values for all the other changes were greater than 0.05, and hence are not significant.

B. Rats

1. Hagan, Fitzhugh et al. (173) studied the oral subacute toxicity of ethyl formate in the rat in an investigation involving a large number of food flavorings and compounds of related structure.

Twenty weanling Osborne-Mendel rats (10 males and 10 females), litter mates, randomized by weight (same weight at each level), were fed diets containing ethyl formate at levels of 1000, 2500, and 10,000 ppm for a period of 17 weeks. A control group of the same number was maintained on the basic diet alone.

Food intake, body weight, and general condition were recorded each week. Fatalities were autopsied, tissues examined for gross pathologic changes, and suspected cause of death recorded; the vital organs were preserved for microscopic study, if not too far autolyzed. Hematologic studies (white cell counts, red cell counts, hemoglobin, and hematocrits) were made at the end of the study. The animals were then sacrificed and all tissues examined macroscopically. Also, liver, kidneys, spleen, heart, and testes were weighed and preserved in 10% buffered formalin-saline, along with the remaining abdominal and thoracic viscera, for histopathologic examination. A detailed microscopic examination of the tissues of six to eight rats (same number of each sex) from the 10,000 ppm group was made.

All animals survived the 17-week feeding period without manifestations of toxic effects or growth retardation. Hematologic findings were normal. No gross or microscopic abnormalities were noted in any of the organs or tissue sections studied.

2. Sporn et al. (435) conducted a chronic toxicity study in rats with formic acid administered in the diet or in the drinking water for varying periods of time.

In the first experiment, young white rats, 40-41 grams BW, eight animals per group, were fed a casein diet containing formic acid at levels of 0.5% or 1.0% for a period of five to six weeks to determine effect on growth. The amount of food ingested, weight gain, and protein efficiency were determined. The results are presented in Table 9. Growth was retarded at both levels of formic acid but the authors state that the degree was not significant statistically.

Table 9. Influence of Formic Acid on the Growth of White Rats (435)
Mean Data Per Group

Group	Initial Indicator	Week						Mean Totals
		I	II	III	IV	V	VI	
I. Casein Control 18.2%	Weight 40.1	56.8	79.0	96.3	116.9	131.9	-	360.8 1.39
	Ingestion	49.1	63.6	77.8	85.4	85.0	-	
	Protein Efficiency	1.86	1.91	1.22	1.32	0.96	-	
II. Formic Acid 0.5%	Weight 40.5	58.6	77.6	90.3	105.3	120.9	-	342.9 1.28
	Ingestion	46.0	63.9	69.9	80.0	83.1	-	
	Protein Efficiency	2.16	1.63	0.99	1.33	1.03	-	
III. Formic Acid 1%	Weight 40.6	60.3	80.7	100.6	117.6	131.0	-	367.9 1.36
	Ingestion	48.9	69.0	80.1	83.3	86.6	-	
	Protein Efficiency	2.20	1.62	1.36	1.12	0.90	-	
IV. Casein Control 11.8%	Weight 40.1	53.5	64.1	75.3	93.5	107.0	115.8	416.2 1.54
	Ingestion	53.9	62.0	71.6	81.0	76.4	71.3	
	Protein Efficiency	2.10	1.58	1.32	1.90	1.49	1.04	
V. Formic Acid 0.5%	Weight 40.1	51.3	63.1	73.1	88.2	98.8	107.1	392.6 1.44
	Ingestion	51.6	64.8	65.4	74.5	70.3	66.0	
	Protein Efficiency	1.83	1.54	1.29	1.71	1.27	1.56	
VI. Formic Acid 1%	Weight 40.1	51.1	63.8	75.9	88.8	100.0	108.3	396.0 1.45
	Ingestion	51.6	65.5	71.4	74.4	69.0	64.1	
	Protein Efficiency	1.82	1.64	1.43	1.46	1.37	1.09	

The second experiment was performed with adult white rats, 179-183 grams body weight, given formic acid in the drinking water at concentrations of 0.5% or 1.0% during fasting and subsequent restoration period of seven days. The effect on weight of vital organs, liver nitrogen and lipid, ascorbic acid content of the suprarenal glands, hemoglobin, and blood catalase coefficient were determined. The results are presented in Tables 10 and 11.

The authors concluded that liver nitrogen values and suprarenal gland weights were significantly lower in animals receiving 0.5% formic acid in comparison with the controls.

In the third experiment, the effect of prolonged administration of formic acid in the drinking water (1.0%) on reproduction, hematology, liver nitrogen, and ascorbic acid content of the suprarenal glands was studied. Sixty adult white rats (10 males and 50 females) were divided into five groups as follows:

- Group I - Untreated controls
- Group II - Females. Formic acid throughout
- Group III - Males. Formic acid throughout
- Group IV - Males and females. Formic acid throughout
- Group V - Females. Formic acid during lactation period only.

Values for the various parameters studied were determined one month, three months, and seven months after placing the animals on the compound. The data are given in Table 12.

The authors state that the most significant finding in this study was the adverse effect of formic acid on survival of offspring during the first seven days of life. Also, hyperchromic anemia, a leucocytosis, a basophile neutropenia, and a slight lymphocytosis, were found in the formic acid groups.

3. Sollmann (430) gave rats, 200-300 gm. body weight, three to six per group, (strain, age, sex, not specified), in the drinking water, formic acid at levels of 8.2, 10.25, 90, 160, and 360 mg/kg BW/day for a period of two to 27 weeks. The animals were fed a standard diet but the treated water was the only source of fluid. Food consumption, fluid intake, and growth rates were determined. The animals were observed also for toxic symptoms throughout the experimental period. The results are presented in Tables 13, 14, and 15.

Table 10. Influence of Formic Acid on Weight of Liver, Kidney, Suprarenal Gland, and Spleen (435)

Mean data per Group				
Group	Liver	Kidney	Suprarenal	Spleen
I. Casein control, 18.2%	5.443	1.302	0.2494	0.424
II. Formic Acid, 0.5%	5.143	1.191	0.2394	0.355
III. Formic Acid, 1%	5.090	1.241	0.2745	0.443
IV. Casein Control, 11.8%	4.318	1.030	0.247	0.310
V. Formic Acid, 0.5%	3.995	1.015	0.226	0.261
VI. Formic Acid, 1%	4.092	1.025	0.226	0.341

Table 11. Influence of Formic Acid on Nitrogen and Lipid Content of the Liver, on Ascorbic Acid Content of Suprarenal Glands, and on the Catalasic Coefficient (435)

Mean Data Per Group				
Group	mg Liver N % Body Wt.	Liver Lipids	Ascorbic Acid Suprarenal Glands % Body Wt.	Catalasic Coefficient
I. Casein control, 18.2%	131.0	0.296	118.8	16.7
II. Formic Acid, 0.5%	128.0	0.228	125.5	19.5
III. Formic Acid, 1%	118.0	0.226	117.5	18.8
IV. Casein Control, 11.8%	115.0	0.214	126.3	17.4
V. Formic Acid, 0.5%	117.0	0.214	135.0	17.5
VI. Formic Acid, 1%	117.0	0.204	132.8	18.8

Table 12. The Influence of Prolonged Administration of Formic Acid on the Reproduction of White Rats. Mean Data Per Group. (435)

Group	Pregnant Females	Offspring Born	Offspring Born Alive	Offspring Alive at 7 days	Offspring Alive at 21 days
Repetition I					
I	6	6.6	6.3	5.0	5.0
II	6	5.5	5.0	0.5	0.5
III	8	6.3	6.1	3.8	3.8
IV	5	6.0	5.4	2.2	2.2
V	6	7.3	7.0	5.5	5.5
Repetition II					
I	6	6.0	4.0	4.0	4.0
II	6	4.1	2.8	2.1	2.1
III	6	6.3	6.0	6.0	6.0
IV	3	4.0	1.0	0.0	0.0
V	7	6.3	5.7	4.7	4.7

No fatalities occurred at the highest level of the acid given (360 mg/kg BW/day) and the author concluded that the occasional deaths at lower concentrations were coincidences. Untreated controls were not included in the experiment.

Food consumption and growth were markedly reduced by formic acid at a level of 360 mg/kg/BW/day for nine weeks after 17 weeks at 90 mg/kg BW/day. Fluid consumption, however, was not depressed. The lower levels of treatment had no adverse effect.

Table 13. Dosage of Formic Acid (430)

CONCENTRATION OF ACID	EXPERIMENT NUMBER	NUMBER OF ANIMALS IN EXPERIMENT	DURATION	MEAN DOSAGE, MILLIGRAMS OF ACID PER KILOGRAM OF RAT PER DAY
Formic acid				
0.01	57	6	11	8.2 (5.6-12.1)
0.01	108	3	14	10.25 (8.4-14.3)
0.1	64	6	15	90.0 (80.0-131.0)
0.25 (after 12 weeks of 0.01)	5796	4	15	160.0 (120.0-210.0)
0.5 (after 17 weeks of 0.1)	6406	3	9	360.0 (290.0-670.0)

Table 14. Effects of Formic Acid on Growth (430)

DRUG AND CONCENTRATION	DOSAGE, MILLIGRAMS PER KILOGRAM OF RAT PER DAY	EXPERIMENT NUMBER	DURATION	OBSERVED WEIGHT	NORMAL WEIGHT	DIFFERENCE IN GRAMS	DIFFERENCE IN PER CENT OF NORMAL WEIGHT	DIFFERENCE PER CENT PER WEEK
Formic acid, 0.01 per cent	8.2	57	10	199	190	+9	+4.6	-0.46
			11	162	192	-30	-15.0	-1.3
Formic acid, 0.01 per cent	10.25	108	14	160	177	-17	-9.0	-0.63
Formic acid, 0.1 per cent	90.0	64	15	275	282	-7	-2.0	-0.13
Formic acid, 0.25 per cent	160.0	5796	2	165	168	-3	-1.8	-0.9
(after 12 weeks of 0.01 per cent)			4	222	175	+47	+26.0	+6.4
			10	235	191	+44	+23.0	+2.3
			15	216	197	+19	+9.0	+0.6
Formic acid, 0.5 per cent								
(after 17 weeks of 0.1 per cent)	360.0	6406	9	180	295	-115	-56.0	-6.7

Table 15. Mortality (430)

DRUG AND CONCENTRATION	DOSAGE	EXPERIMENT NUMBER	NUMBER OF ANIMALS	WEEKS OF FATALITIES	TOTAL DURATION	PER CENT OF FATALITIES
Formic acid, 0.01 per cent	8.2	57	6	11	11	16
Formic acid, 0.01 per cent	10.25	108	3	0	14	0
Formic acid, 0.1 per cent	90.0	64	6	0	15	0
Formic acid, 0.25 per cent (after 12 weeks of 0.01 per cent)	160.0	5796	4	2	15	25
Formic acid, 0.5 per cent (after 17 weeks of 0.1 per cent)	360.0	6406	3	0	9	0

C. Guinea Pigs

Amdur (007) exposed guinea pigs, seven to 16 per group, (strain, age, sex, and weight not given), to air containing formic acid at levels ranging from 0.34 to 42.5 ppm, alone and with sodium chloride aerosol, for a period of one hour. Intrapleural pressure, tidal volume, and the rate of flow of gas in and out of the respiratory system were determined at five-minute intervals to assess the effect of the irritant. Normal values were obtained during a pre-treatment control period of one-half hour. Each animal, therefore, served as its own control. Also studied were responses to the irritant administered through a tracheal canula to by-pass the protective effect of the upper airway. Post-exposure periods varied from one to three hours. The results are presented in Tables 16 and 17, and Figs. 4, 5 and 6.

Formic acid at concentrations as low as 0.3 ppm for one hour caused an increase in resistance (pulmonary flow) and a decrease in compliance (elastic recoil) which were statistically significant. Concentrations of 1-3 ppm produced an increase in respiratory work, and above 10 ppm, respiration rate and minute volume decreased significantly. The effect of formic acid was not potentiated by the sodium chloride aerosol. In this study formic acid proved to be a more potent respiratory irritant than formaldehyde.

D. Rabbits

1. Smyth et al. (429) subjected male New Zealand giant albino rabbits, 2.5-3.5 kg BW, four per group, to graded doses of ethyl formate applied to the skin (hair removed by clipping) over about 1/10 of the body surface and retained in contact with the skin by means of an impervious plastic film. After an exposure period of 24 hours, the film was removed and the animals observed for a period of 14 days for signs of systemic toxic effects. All rabbits survived the maximum dosage, 20 mg/kg BW, that could be retained in contact with the skin by the method employed. No adverse effects were reported.

The same authors (429) also applied varying amounts and dilutions of ethyl formate to the clipped skin and to the eyes of albino rabbits, five per group (strain, age, weight, and sex not given), and determined the degree of primary skin irritation or eye injury over a period of 24 hours. The results were scored on the basis of a "10-grade ordinal series" in which Grade 1 indicated the mildest effect and Grade 10 the most severe reaction. Ethyl formate produced a so-called severe corneal burn (Grade 4) from a relatively small amount introduced into the eye; dermally, it was quite non-toxic (Grade 1).

Table 16. Response of Guinea Pigs to One-Hour Exposure to Formic Acid (007)

HCOOH-P.P.M.		0.34±0.02	1.0±0.08	2.8±0.15	6.6±0.5	13.5±0.9	42.5±3.8
Number of Animals		7	8	8	16	8	7
Resistance cm H ₂ O/ml/sec	Control	0.78	0.71	0.77	0.77	0.81	0.86
	Exposure	1.01 ^b	1.02 ^b	1.28 ^b	1.40 ^a	1.56 ^a	2.08 ^a
Exposure-Control		+0.23	+0.31	+0.51	+0.63	+0.75	+1.22
St-Standard Error of Difference		0.053	0.059	0.165	0.049	0.100	0.157
Compliance ml/cm H ₂ O	Control	0.26	0.23	0.24	0.25	0.21	0.20
	Exposure	0.21 ^d	0.18 ^b	0.19 ^c	0.19 ^a	0.14 ^b	0.13 ^b
Tidal Volume ml	Control	1.85	1.91	1.88	1.84	1.70	1.70
	Exposure	1.83	1.71	2.12	1.82	1.82	1.92
Frequency Breaths/min	Control	95	96	101	100	107	100
	Exposure	89	101	90	99	98	58 ^b
Minute Volume ml/min	Control	176	183	183	184	183	170
	Exposure	165	172	175	179	173	110 ^b
Elastic Work g-cm/min	Control	624	762	744	677	736	722
	Exposure	710	820	1065 ^c	974 ^c	1160 ^c	822
Resistive Work g-cm/min	Control	990	980	1140	1070	1100	1020
	Exposure	1100	1250	1917 ^b	1860	2040 ^b	1060
Total Work g-cm/min	Control	1614	1742	1884	1747	1836	1742
	Exposure	1810	2070	2982 ^b	2834 ^c	3200 ^b	1882

Different from control at $P < a = .001, b = .01, c = .02, d = .05$

Table 17. Resistance Values One Hour After End of Exposure (007)

HCHO							
HCHO-P.P.M.	0.05	0.31	0.58	1.22	3.5	11.0	49
Number of Animals	18	13	23	4	10	10	8
Resistance control	0.71	0.64	0.64	0.80	0.68	0.62	0.65
cm H ₂ O/ml/sec 1 hr. post exposure	0.82	0.62	0.71	0.93	0.82	0.75	1.60
\bar{x} -Post Exposure--control	+0.11	-0.02	+0.07	+0.13	+0.14	+0.13	+0.37
S \bar{x} -Standard Error of Difference	0.05	0.056	0.038	0.125	0.080	0.071	0.065
Level of Significance $P <$	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	0.01
HCHO--Tracheal Cannula							
HCHO-P.P.M.	0.90	5.2	20	50		1.14	3.6
NaCl-mg/m ³						10	10
Number of Animals	7	7	10	6		8	6
Resistance control	0.40	0.31	0.44	0.48		0.38	0.44
cm H ₂ O/ml/sec 1 hr. post exposure	0.46	0.41	0.87	1.04		0.98	1.24
\bar{x} -Post Exposure--control	+0.06	+0.10	+0.34	+0.56		+0.60	+0.80
S \bar{x} -Standard Error of Difference	0.036	0.049	0.213	0.198		0.207	0.196
Level of Significance $P <$	N.S.	N.S.	N.S.	0.05		0.05	0.01
HCOOH							
HCOOH-P.P.M.	0.34	1.0	2.8	6.6	13.5	42.5	
Number of Animals	7	8	8	16	8	7	
Resistance control	0.78	0.71	0.77	0.77	0.81	0.86	
cm H ₂ O/ml/sec 1 hr. post exposure	0.75	0.94	1.02	1.05	0.98	1.54	
\bar{x} -Post Exposure--control	-0.03	+0.23	+0.25	+0.38	+0.17	+0.68	
S \bar{x} -Standard Error of Difference	0.090	0.154	0.146	0.077	0.096	0.222	
Level of Significance $P <$	N.S.	N.S.	N.S.	0.01	N.S.	0.02	




Figure 4. Time Course of the Resistance Change Produced by Formic Acid (007)




Figure 5. Comparison of Changes in Resistance and Compliance Produced by Equal Concentrations of Formaldehyde and Formic Acid (007)




Figure 6. Comparison of the Changes in Respiratory Work Produced by Equal Concentrations of Formaldehyde and Formic Acid (007)

2. Lund (272) in a study on the metabolism and excretion of methanol and formic acid, injected two full-grown male rabbits, weighing 3150 and 3300 grams, subcutaneously with a single dose of sodium formate, consisting of 317 and 303 mg/kg BW, respectively. Although the experimental period lasted only 48 hours, apparently neither animal manifested any toxic symptoms.

3. Sollmann (430) reported that an early study by Lebbin (1916) revealed that continuous administration of either acetic acid or formic acid (dosage, route, and duration not given) caused kidney damage.

4. Gross (260) noted that rabbits, like guinea pigs, suffered marked depression of the central nervous system when placed in an atmosphere containing ethyl formate at concentrations of 40 to 130 mg per liter, and that pneumonia frequently developed. Subcutaneously, however, doses as large as 1000 mg/kg BW were tolerated without local or systemic symptoms.

5. von Oettingen (491), in a review of early studies on the effect of ethyl formate administered intravenously to the rabbit, noted that intrajugular injection of 28 mg/kg BW of the undiluted ester caused an increase of the respiratory volume without depressing the CNS, whereas, in another study, doses as high as 250 mg/kg BW intravenously (anatomic site not given) in the form of a 5% solution, were tolerated without evidence of any effect.

E. Dogs

Lund (271) in a study of the metabolism and excretion of methanol and formic acid, injected a full-grown male dog, 10.1 kg BW, (strain, sex, and age not given) subcutaneously with a single dose of sodium formate, consisting of approximately 200 mg/kg BW (40 ml of a 5% solution). The animal manifested no toxic signs over an experimental period of 27 hours.

In a second experiment by the same author, a single dose of approximately 100 mg/kg BW (50 ml of a 2% solution), injected into the washed, empty bladder through a catheter tube, was tolerated without toxic symptoms.

F. Sheep

Neumark et al. (315) gave (stomach tube or rumen canula) two clun forest wethers (age and weight not specified), single doses of formic acid, approximately 150 mg/kg BW. This was done in an effort to determine if the appetite depressant effect of this compound in sheep was due to systemic action or to

a local effect on one or more areas of the stomach. (Previous studies showed that the quantity of silage eaten by sheep was reduced when fermentation products, such as formic acid, formaldehyde, and histamine were present). In another experiment, the acid was injected intravenously. Also, in one test, sodium formate was used in place of formic acid. The effects of histamine, and of histamine in combination with formic acid, were determined in other experiments. Control data (no treatment) were obtained on all of the sheep in preliminary experiments. During the experimental period, the animals were maintained on a ration consisting of 350 grams of chopped hay with 100 grams of flaked maize fed twice daily. The effect on eating time of offered food was determined. The results are presented in Tables 18, 19, 20 and 21.

The data obtained showed that the area between the cardia and fundic portions of the abomasum is sensitive to formic acid. When the compound was introduced into this area at a level of 150 mg/kg BW, there was a marked adverse effect on the eating time of one of the two animals used (sheep 174). A combination of formic acid and histamine had a much stronger depressant effect (see Table 18). Experiments with dyes revealed that substances introduced into the above-mentioned area of the stomach passed rapidly into the abomasum. Formic acid alone, or in combination with histamine, introduced into the rumen via a canula, had no effect on the time sheep took to eat offered food. It is known that formic acid is rapidly destroyed in the rumen and provides a source of hydrogen for methane formation. Administration of formate intravenously had no effect on the eating behavior.

The authors suggest that the depressing effect of formic acid on appetite is caused by its local irritant action on the nerve endings of the stomach as it rapidly penetrates the epithelial layer. Sodium formate does not cause this effect.

G. Cattle

1. Castle (074) fed 12 Ayrshire cows, (age and weight not given), grass silage containing formic acid at a level of 1/2 gallon per ton of herbage, for a period of twelve weeks. Control groups on silage without additive (wilted and unwilted) were included. The silages were fed to the cows ad libitum with a supplement of moist barley and groundnut cake. Food intake, weight gain, milk yield, and solids-not-fat content of the milk were determined.

No toxic effects were noted at any time during the study. The formic acid-treated silage proved to be superior to the control diet with respect to all parameters.

Table 18. The Effects of Introducing Formic Acid, Histamine, Saline and Other Agents into the Rumen via a Stomach Tube on the "eating time" of Sheep (315)

No. of trials	Solution introduced	Effect on 'eating time'
5	1 l. saline	None
4	1 l. 0.2 M formic acid	Marked effect in 3 trials. Slight effect in 1 trial
4	1 l. 0.2 M formic acid + 360-541 mg histamine (3 trials) or 902 mg histamine (1 trial)	None at lower rate of dosage. Marked at high rate
Uncontrolled length of tube used		
7	1 l. 0.2 M formic acid + 497-660 mg histamine (tube introduced to 72-73 cm)	Marked effect in all trials
1	1 l. 0.2 M formic acid plus 497 mg. histamine (tube introduced to 37 cm)	None
1	1 l. 0.2 M acetic acid (tube introduced to 72 cm)	None
1	1 l. 0.25 M acetic acid plus 497 mg. histamine (tube introduced to 72 cm)	Marked effect
2	1 l. 0.2 M sodium formate (tube introduced to 72 cm)	None

Table 19. The Responses of Sheep to Infusions of Formic Acid, Histamine or Both into the Abomasum (315)

Sheep	Treatment	Response
(Infused into pyloric region)		
174	0.016 M formic acid plus 42 mg histamine	None
174	0.026 M formic acid + 75 mg histamine	Rapid respiration. Finished food in 31 min.
174	94.5 mg histamine	Rapid respiration. Finished food in 35 min.
174	0.08 M formic acid	None
174	0.19 M formic acid	None
(Infused into fundic region)		
265	0.15 M formic acid plus 120.6 mg histamine	Normal respiration. Stopped feeding after 6 min. 150 g left after 42 min. Feeding normally after 24 min.
70		Normal respiration. Stopped eating after 9 min. and did not feed normally for 2 days
34		Normal respiration. Did not feed after infusion and food consumption depressed for several days

Table 20. The Response of Food Intake to Infusion of Formic Acid and/or Histamine into the Jugular Vein (315)

No. of trial	Treatment*	Response
1	0.0005 M formic acid plus 1 mg histamine	Finished eating after 27 min; several very slight interruptions during eating
2	0.001 M formic acid	No
3	0.001 M formic acid 1 mg histamine	No
4	0.002 M formic acid 2 mg histamine	Respiration rate raised, eating speed lowered (31 min)
5	0.004 M formic acid 1 mg histamine	No
6	0.004 M formic acid	No (finished eating after 24 min.)
7	3.2 mg histamine	Respiration rate raised, finished eating after 32 min.

* Formic acid and/or histamine were dissolved in 10 c.c. and in 8 c.c. saline in trials 1-6 and 7 respectively

Table 21. The Concentration of Acetic and Formic Acids in the Jugular Blood of Sheep no. 174 Before and After Treatment* (315)

No. of trial	Time of blood collection	Infusion by stomach tube		Infusion by rumen fistula	
		Formic acid ($\mu\text{g/ml}$ blood)	Acetic acid ($\mu\text{g/ml}$ blood)	Formic acid ($\mu\text{g/ml}$ blood)	Acetic acid ($\mu\text{g/ml}$ blood)
1	Before treatment	14.0	43.7	5.9	29.0
2	6 min after treatment	---	---	16.9	27.0
3	30 min after treatment	76.4	68.7	8.9	28.7

* One litre 0.2 M formic acid solution, containing 861 mg histamine, was infused by stomach tube into the rumen of sheep no. 174, and 5 days later an equal solution, containing only 581 mg histamine, was introduced into the rumen of the sheep by rumen fistula.

2. Waldo et al (496) fed Holstein heifers, paired on body weight (89-337 kg), Williamsburg alfalfa and Potomac orchard grass silage, treated at the field chopper with an additive of 0.49, 0.52, or 0.48 (w/w) of 90% formic acid solution. The silages were stored for 28, 29, and 128 days, respectively, before feeding (experiments 1, 2, 4). Each silage was fed to the heifers ad libitum as the sole ration. Control silages were included in each experiment. The organic acid concentration is shown in Table 22. Food intake, digestibility, and growth rates were determined. The results are presented in Tables 23 - 26. Another experiment (Experiment 3) was conducted in which the silage was treated with a combination of formic and acetic acids.

No toxic effects were reported. Weight gains of the animals on the formic acid silage were better for the most part than those of animals on the control ration.

3. Bovine erythrocytes are hemolyzed by ethyl formate in vitro. The minimal lytic concentration found was 5.25 grams (0.71 mole) per liter (491).

Table 22. Organic Acid Concentration in Silages. (496)

Criteria	Experiment 1		Experiment 2		Experiment 3		Experiment 4		Overall means		Standard error of mean ^a
	Formic	Un-treated	Formic	Un-treated	Formic	Un-treated	Formic	Un-treated	Formic	Un-treated	
n samples	4	4	4	4	4	4	5	5
acetic (% DM)	5.29	7.42	2.96	4.14	1.06	1.62	5.57	0.31	3.72	3.35	1.21g
butyric (% DM)	4.94	4.13	3.79	1.52	.70	.73	1.22	2.57	2.68	2.24	.35g
propionic (% DM)	.08	.02	.16	.07	.01	.07	.06	.76	.08	.23	.14g
lactic (% DM)	.06	0	.17	.06	0	.04	.13	1.85	.08	.59	.33g
malic (% DM)	.56	.10	.28	.18	.79	.03	.99	.15	.66	.13	.12g
total (% DM)	10.96	11.67	7.35	5.98	2.56	2.54	7.97	5.36	7.21	6.16	.19g

^a Standard error of a mean and statistical significance is for overall means: e, $P < .05$ and g, $P > .10$.

Table 23. Least Squares Treatment within Experiment Means and Standard Errors of Ad Libitum Forage Intake and Heifer Growth (496)

Criteria	Experiment 1		Experiment 2		Experiment 3		Experiment 4		SE of a mean ^b
	Formic	Un-treated	Formic	Un-treated	Formic	Un-treated	Formic	Un-treated	
Days	72	72	69	69	84	84	36	36	..
No. animals	15	15	14	14	10	10	10	10	..
Daily gain (g)	763	653	570	651	931	824	949	61	49.0a
Daily gain (g/kg ^{0.75})	12.7	10.8	9.6	11.0	13.5	11.7	12.4	.9	.74a
Daily DM consumed (% BW)	2.53	2.06	2.35	2.54	2.57	2.04	1.80	1.46	.054a
Daily Meal DE consumed	18.5	18.9	16.2	16.9	18.6	19.6	16.0	12.0	.41a
Daily feed DE/kg ^{0.75}	300	311	276	286	263	276	209	160	.7a
Daily Meal DE for growth	9.7	10.2	7.7	8.3	8.3	9.3	4.9	1.1	.35a
Meal DE for growth/kg gain	12.5	15.5	11.8	13.4	8.7	11.6	5.1	6.6	.4g
Gain/retention DM stored (kg)	98	100	84	96	122	82	146	11	

^a Standard error of a mean is for n = 10. Statistical significance is for experiment by treatment interaction: a, P < .001 and g, P > .10.

Table 24. Least Squares Treatment Means and Standard Errors of Ad Libitum Forage Intake and Heifer Growth (496)

Criteria	Formic	Untreated	SE of a mean ^b
No. animals	49	49	..
Daily gain (g)	808	548	142.4g
Daily gain (g/kg ^{0.75})	12.1	8.6	1.88g
Daily DM consumed (% BW)	2.32	2.33	.076g
Daily Meal DE consumed	17.3	16.8	.77
Daily feed DE/kg ^{0.75}	262	258	10.1g
Daily Meal for growth	7.6	7.2	.71g
Meal DE for growth/kg gain	10.4	11.8	.76g
Gain/retention DM stored (kg)	113	72	23.7g

^b Statistical significance: g, P > .10.

Table 25. Least Squares Treatment Within Experiment Means and Standard Errors for Digestibility and Nitrogen Utilization (496)

	n	Dry matter	Energy	Cellulose	Hemi- cellulose	Lignin	Nitrogen digested	Nitrogen retained	Fecal solubles
									(g/100 g feed DM)
Experiment 1		(%)					(g/day)		
Formic ad lib	4	62.2	63.9	61.8	50.1	5.5	71.3	41.7	16.1
Formic maintenance	4	64.2	65.6	66.5	50.8	8.8	72.1	10.6	14.9
Untreated ad lib	4	64.5	65.9	65.8	35.6	14.9	72.6	46.2	15.8
Untreated maintenance	4	65.8	67.5	66.9	41.8	11.7	75.2	12.0	14.8
Experiment 2									
Formic ad lib	4	62.5	62.5	70.5	61.9	16.6	70.4	33.0	16.2
Formic maintenance	4	63.8	63.4	70.8	66.2	12.8	71.4	9.7	15.1
Untreated ad lib	4	63.2	62.7	69.1	60.8	17.1	66.8	20.9	17.0
Untreated maintenance	4	65.2	64.4	71.0	67.0	18.7	68.5	21.8	16.1
Experiment 3									
Formic ad lib	4	56.0	55.0	60.6	62.4	6.5	58.4	26.1	21.6
Formic maintenance	4	57.0	55.7	60.2	57.6	3.9	61.4	2.8	19.4
Untreated ad lib	4	58.2	56.4	65.6	44.6	16.6	57.4	29.8	20.0
Untreated maintenance	4	58.4	57.6	66.4	51.4	16.6	59.4	6.5	20.4
Experiment 4									
Formic ad lib	4	56.6	57.0	67.2	73.6	16.0	61.9	30.8	20.8
Formic maintenance	4	56.8	57.5	67.3	74.0	11.2	63.0	14.8	20.4
Untreated ad lib	4	46.6	50.7	58.2	62.8	9.7	51.8	16.6	24.4
Untreated maintenance	4	48.3	51.3	59.5	67.1	10.5	53.8	17.0	23.6
SE of a mean ^b		1.31b	1.47c	1.74b	3.26g	5.78d	1.70d	7.08b	.56b

^b Statistical significance is for experiment by treatment interaction: b, P < .005; d, P < .025 and g, P > .10.

Table 26. Least Squares Treatment Means and Standard Error for Digestibility and Nitrogen Utilization (496)

Criteria	Formic	Un-treated	SE of a mean ^a
n	32	32	
Dry matter (%)	59.9	58.8	1.12g
Energy (%)	60.1	59.6	.79g
Cellulose (%)	66.0	65.3	1.19g
Hemicellulose (%)	62.1	53.9	1.27b
Lignin (%)	10.5	9.4	3.40g
Nitrogen (% digested)	66.2	63.2	1.02f
Nitrogen (g retained/day)	21.2	12.9	5.36g
Fecal solubles (g/100 g feed DM)	18.1	19.0	.38g

^a Statistical significance: b, $P < .005$; f, $P < .10$ and g, $P > .10$

H. Human toxicity

Sollmann (430) and von Oettingen (491) have reviewed the early studies on the effect of formic acid and formates on human volunteers. In one study 500 mg. amounts of formic acid in lemonade given daily for four weeks caused no harmful effects. Larger doses (amount not specified) caused local reactions analogous to those produced by acetic acid.

Ingestion of 3-4 grams of sodium formate (50-70 mg/kg BW) caused albuminuria and hematuria in one of three human subjects but the condition cleared up after five days (491). In another study, daily doses of 150 mg/kg BW per os for some time (duration not given) caused no harm (430). Intravenously, a single dose of four grams (20 ml of a 20% soln.) of sodium formate was tolerated without evidence of toxicity (491). On the other hand, Stern (445) found that large doses, 2-3 grams given several times daily, may cause vertigo, nausea, vomiting, albuminuria, hematuria, tenesmus, dyspnea, and lowered body temperature.

Formic acid and formates are quite toxic for man via the respiratory tract. Exposure to the vapors or dust may cause pallor, hyperemia of the conjunctivae and sclera, and gastrointestinal distress. Inhalation in concentrations as low as 32 mg per liter of air causes progressive irritation of the eye and mucous membranes which may last four hours after withdrawal of the

irritant (491). The maximum acceptable concentration of ethyl formate, according to the American Conference of Governmental Industrial Hygienists, in air is 100 ppm for 8-hour exposures daily (009).

Gray (162) studied the effect of a series of organic acids on human tooth enamel by subjecting sections of teeth to varying concentrations of the acids over a pH range of 3.0 to 6.0 and determined the degree of caries-like lesion formation. The results revealed that formic acid at pH 4.5 had a pronounced effect on tooth enamel. At pH 4.0, the effect was considerably greater (see Figs. 7 and 8).

Formic acid is very corrosive to the skin and mucous membranes, and precautions should be taken to avoid direct contact with the chemical or breathing formate dusts and mists (438). Even though skin contact may be slight and only moderately painful at first, the after-effects can be severe. Following the initial reaction of hyperemia and erythema, the skin may show "a cooked appearance, and after 24 hours, parchment-like necroses which are slow to heal". Orally, formic acid and formates cause irritation of the gastrointestinal tract followed by vomiting and diarrhea (491).

A number of conflicting reports have appeared concerning possible roles of formic acid and formaldehyde, which are intermediate oxidation metabolites, in methyl alcohol poisoning in man. Lund (270) reported on concentrations of methanol and formic acid in the blood and urine of human victims poisoned with methyl alcohol. The results are presented in Tables 27 and 28.

Table 27. Formic Acid Excretion of Human Subjects Poisoned with Methanol. (270)

A: Man, aged 31, weight 62 kg, consumed 25 ml (20 g) methanol (Pohl 1893).

B: Woman, aged 25, consumed about 80 g methanol (Harrop and Benedict 1920)

Days after intake	excreted formic acid in urine			
	mg A	in 100 ml B	mg A	total B
1	2.3	--	37.4	--
2	32.5	126	420	--
3	6.1	--	89.6	--
4	2.4	60	29.2	--
11	--	7	--	--




Figure 7. The enamel dissolution rate during caries-like formation in 6% hydroxyethylcellulose + acid solutions is shown for four acids with decreasing dissociation constants. The solid lines are calculated values from a single equation fitted to the results. (162)




Figure 8. The enamel dissolution rate as a function of pH and total available acid is shown for the 4 different acids of Fig. 7. The rate is pH dependent and linear with respect to total acid concentration except for acetate at high pH levels. (162)

Table 28. The Concentration of Methanol and Formic Acid in Blood, Liver and Urine from Human Beings Dying of Methanol Poisoning (270).

Case No.	mg methanol in 100 g of			mg formic acid in 100 g of		
	blood	liver	urine	blood	liver	urine
10. V.H.	74	106	140	68.1	59.6	785
11 O.R.*		--	240	--	98.6	475

Watts (504) studied the effect of formic acid, formaldehyde, and methanol on the oxidation of glucose, succinate, and ascorbate by rat liver slices and homogenates, in order to shed some light on the mechanism of methanol poisoning (see this section, A.). Although formic acid, in concentrations approximating those found in fatal methanol poisoning in the rabbit, depressed respiration by 50%, formaldehyde was found to be ten times more potent in this respect. On the basis of a comprehensive review of the literature, plus his own experimental findings, Sammartino (403) concluded that methanol per se played the main role in methyl alcohol poisoning in man. The known physiologic effects of these compounds, modes of action, and pattern of toxic effects in methanol poisoning led him to this view.

III. Long-term Studies

A. Rats

Malorny (275) recently concluded three long-term oral toxicity studies on the rat with calcium formate and sodium formate administered in the drinking water.

The first experiment was started with Wistar rats; eight males and twenty-four females, 160-210 grams body weight. These animals were given calcium formate in the drinking water at a level of 0.2% throughout their life span. Fluid intake measurements indicated a dosage of 150-200 mg/kg BW daily. A control group of eight animals of the same specifications was included. The experiment was continued with the progeny of the original animals. After more than three years and five generations, no deaths or toxic symptoms attributable to the compound were noted. Fertility, pregnancy, and development of fetuses and offspring also were normal. The data are presented in Tables 29 and 30.

Table 29. Effect on Rats of Calcium Formate (0.2%) Administered^a Orally in the Drinking Water (275)

Weeks	0.2% Ca(HCOO) ₂ in the drinking water				Controls	
	I♀	II♀	III♀	IV♂	V♀	VI♂
1	175 +8.9	170 +12.5	170 +11.0	195 +9.4	170 +11.2	200 +9.8
2	205	190	185	230	195	235
3	220	190	195	250	210	260
4	230	200	200	270	225	275
5	240	205	210	285	235	288
6	240	215	220	300	238	290
7	240	220	225	310	240	305
8	255	220	230	320	260	318
10	262	228	235	340	268	340
12	275 +10.6	240 +9.8	238 +9.5	353 +13.4	270 +9.0	360 +14.2
15	283 +9.5	245 +10.5	240 +13.0	355 +11.8	290 +10.6	370 +12.7

^aFour rats per group. Body weights - Mean values and Standard Deviations

Table 30. Effect on Rats of Generations I-III of Calcium Formate (0.2%) Administered^a Orally in the the Drinking Water (275)

	I. Generation		II. Generation		III. Generation	Control
	n=11	n=10	n=9	n=8	n=10	n=10
Weight at birth	5.0 +4.6	5.8 +6.1	5.5 +3.3	4.9 +5.5	5.6 +5.8	5.2 +5.2
1 week	16.0	19.5	18.2	15.9	14.8	14.5
2 week	34.0	32.5	31.7	28.0	29.1	33.0
3 week	49.0	55.0	48.0	48.0	52.0	53.0
4 week	79.0 +3.8	84.0 +3.7	78.0 +4.0	85.0 +3.2	76.0 +3.6 ¹⁾	83.0 +4.2
5 week	108	118	112	109		113
6 week	128	138	136	134		145
7 week	160	159	155	152		165
8 week	179 +3.9	174 +4.8	177 +6.0	180 +4.3		182 +5.1
10 week	216	220	217	224		221
14 week	291	297	301	295		293
18 week	297	309	-	-		-
20 week	309 +5.0	318 +6.2	310 +6.8	313 +5.2		320 +5.8

¹⁾The test is not pursued further.

^aMean Body Weights with Standard Deviation

The second experiment with calcium formate by Malorny (275) was similar in all respects to the first except that the formate concentration was doubled (0.4%) in the drinking water, and the data reported were for two generations of animals over a period of two years. Again, no deaths or toxic effects resulted from the calcium formate administered. Moreover, gross and pathologic examination of more than 250 rats from three generations employed in these experiments, revealed no evidence of chronic poisoning or malignancy.

The third experiment in this series conducted by Malorny (275) was started with the same number of Wistar rats (375 grams mean body weight) as used in the preceding experiments. Sodium formate, however, was used in place of calcium formate, and the concentration in the drinking water was increased to 1%. The daily intake was found to be 274 mg per animal (730 mg/kg BW, approx.). Physiologic studies revealed that 13.8% of the compound ingested, calculated as formic acid, was excreted in the urine daily. No fatalities or toxic effects attributable to the compound over the experimental period of 1-1/2 years were reported.

IV. Special Studies

A. Bacteria

1. Demerec (097) investigated formic acid and a number of other compounds for mutagenic potentiality on the basis of their ability to induce back-mutation from streptomycin-dependence to non-dependence in Escherichia coli (strain Sd-4). Varying concentrations of formic acid in distilled water (See Table 31) were inoculated with streptomycin-dependent E. coli (Sd-4), and incubated at 37°C for six hours. Untreated control cultures were prepared in an analogous manner by inoculating an equal volume of distilled water with the test organism. At the end of the exposure period, both treated and control cultures were plated on streptomycin agar to calculate the percentage of survivors, and on streptomycin-free agar to determine the number of mutants. The results are presented in Tables 31 and 32.

Formic acid was one of 19 compounds found to be mutagenic in this study (Table 32). The data indicate it to be moderately mutagenic, in comparison with the others, with respect to its ability to induce back-mutation in Escherichia coli (Sd-4) from streptomycin-dependence to non-dependence.

Table 31. Results of Experiments Testing Induction of Back-mutations in Streptomycin-dependent *E. coli* Treated with Certain Chemicals (097)

Chemical	Concentration %	Treatment (hours)	Survival %	Treated			Control		
				No. of plates	Total no. of bacteria	Mutants per 10 ⁸ bacteria	No. of plates	Total no. of bacteria	Mutants per 10 ⁸ bacteria
Lactic acid (continued)	0.021	3	7.5	8	1.1 × 10 ⁸	40.2			4.5
	0.020	3	6.8	8	1.7 × 10 ⁸	17.0	5	1.6 × 10 ⁸	15.3
	0.020	3	4.6	8	6.7 × 10 ⁸	32.9	5	9.2 × 10 ⁷	27.3
	0.020	4	2.5	8	2.9 × 10 ⁷	34.7	6	8.5 × 10 ⁷	9.4
	0.020	3	1.9	8	2.6 × 10 ⁷	11.4	6	1.0 × 10 ⁸	4.8
Sodium lactate	2.5	3	85.	8	1.5 × 10 ⁸	7.3	5	1.1 × 10 ⁸	12.1
	3.0	3	61.	8	1.7 × 10 ⁸	16.4	5	1.6 × 10 ⁸	15.3
	2.0	24	8.3	8	8.7 × 10 ⁷	5.8	6	7.9 × 10 ⁷	10.1
	2.5	24	1.4	8	3.8 × 10 ⁷	5.2	5	1.7 × 10 ⁸	9.2
Formic acid	0.0070	3	100.	8	1.5 × 10 ⁸	18.1	6	1.1 × 10 ⁸	2.7
	0.0060	3	93.	8	1.4 × 10 ⁸	18.4			2.7
	0.0050	3	77.	8	1.3 × 10 ⁸	20.2	5	1.1 × 10 ⁸	12.1
	0.0065	3	30.	8	8.2 × 10 ⁸	44.0	5	1.7 × 10 ⁸	9.2
	0.0060	3	20.	8	5.0 × 10 ⁸	43.3	5	1.6 × 10 ⁸	15.3
	0.0065	3	19.	8	4.6 × 10 ⁸	24.4	5	1.5 × 10 ⁸	3.3
	0.0060	3	15.	8	1.4 × 10 ⁸	27.7	6	7.1 × 10 ⁷	2.8
	0.0075	3	3.4	8	8.2 × 10 ⁷	38.0			3.3
	0.0070	3	2.8	8	2.6 × 10 ⁷	38.5			2.8
Formaldehyde	0.01	3	61.	2	7.6 × 10 ⁷	400.	2	1.3 × 10 ⁸	26.2
	0.01	3	58.	5	6.8 × 10 ⁸	133.	4	9.4 × 10 ⁸	15.5
	0.02	3	58.	5	6.8 × 10 ⁸	157.			15.5
	0.01	3	48.	4	3.7 × 10 ⁷	245.	4	7.6 × 10 ⁷	79.
	0.01	3	48.	4	3.7 × 10 ⁸	470.	4	7.6 × 10 ⁸	68.
	0.01	3	21.	8	1.3 × 10 ⁸	120.	4	7.3 × 10 ⁷	12.3
	0.02	3	16.	8	1.0 × 10 ⁸	129.			12.3
	0.04	3	9.0	7	1.5 × 10 ⁸	326.			15.5
	0.05	3	1.7	10	3.2 × 10 ⁸	560.			79.

Table 32. Summary of the Data on Mutagenic Chemicals, Showing Highest and Lowest Frequencies of Mutants Observed in Experiments Using Different Treatments (Expressed in Percentages of Survivors) (097)

	Survivors (per cent)			Controls
	50-100	5-50	less than 5	
No. of mutants per 10 ⁸ bacteria				
Boric acid	27	16-82	19-57	6-31
Ammonia	5	2-10	38-121	3-13
Hydrogen peroxide	20-29	121-124	27-175	5-7
Copper sulfate	6	4-21	40-56	2-21
Acetic acid	8	15-36	57	7-10
Lactic acid	...	11-40	11-35	5-27
Formic acid	18-20	24-44	38-39	3-15
Formaldehyde	133-400	120-326	560	12-79
Phenol	3-85	...	124-820	1-75
Alpha-dinitrophenol	3-34	14-56	6-97	3-13
Trinitrophenol	15	9-132	27-68	3-14
Carbamate, ethyl	19-27	7-53	40-99	6-20
<i>n</i> -propyl	12-28	8-30	10-35	2-12
<i>n</i> -butyl	21	5-21	6-9	2-12
isocetyl	8-11	6	7	2-4
Neutral red, light	57	118-122	178	7-11
dark	11-17	7-15
Acridine	...	59-78	...	14
Caffeine	16	13-35	20	3-18
Necrosin	16-26	18	172	4-10

2. Freese et al (141) reported that formic acid and two other oxidation products of formaldehyde, as well as formaldehyde itself, at levels of 10^{-1} M and 10^{-2} M, did not inactivate or mutate transforming bacterial DNA in vitro at a significant rate in their study.

B. Insects

Stum-Tegethoff (450) investigated the susceptibility of Drosophila germ cells to the mutagenic effect of formic acid, formaldehyde, and some related compounds with respect to sex-linked lethals in males.

A standard strain of Drosophila melanogaster, Oregon-K (inbred for about 10 years), with a spontaneous mutation rate of 0.15% sex-linked lethals in males, was used throughout. Adult males were subjected to the compounds in the vapor phase, and larva to their effect when incorporated in a nutrient medium of agar, killed brewer's yeast, corn meal, and sucrose - pH 4.2-5.6. To test for sex-linked lethals, about 50 treated males were individually crossed with one or two virgin females of the M-5 strain (inbred for about 10 years). Three successive broods were produced by transferring the males to two fresh virgins every third day. The results are presented in Tables 33, 34, and 35.

Formic acid at a level of 0.1% (final concentration), alone and in combination with formaldehyde, proved to be mutagenic for Drosophila germ cells. All germ layers were susceptible in the vapor phase. Spermatocytes I were especially sensitive to their action per os. Volatile acid was necessary for the mutagenic activity of formaldehyde in the vapor phase and a low pH for all compounds in the feeding experiment. When the pH was raised to 7.5, for instance, the induced mutation rate was only slightly higher than the spontaneous rate. The author postulated that the mutation mechanism involved catalase inhibition which promoted the formation of peroxides or free radicals which in turn interfered with DNA replication.

C. Chick embryos

Malorny (275) injected 2 and 4-day old embryonated chicken eggs with single doses of sodium formate (5, 10, or 20 mg per egg) and then observed the embryos for toxic signs and teratogenic effects throughout the hatching period.

Table 33. Formic Acid-induced Mutations in Drosophila melanogaster, Vapor Experiments (450)

treatment		brood							
substance	concentration	1st		2nd		3rd		total	
	%	n ¹	% ²	n	%	n	%	n	%
nF	7.0	—	—	—	—	—	—	3803	0.16
aF	7.0	436	0.69	712	1.97	423	1.18	1571	1.49
FA	0.1	1008	1.12	1139	1.42	850	1.41	3048	1.31
AA	0.1	1703	0.82	1359	0.66	1270	1.10	4332	0.85
HA	0.01	1047	0.67	977	0.61	596	0.50	2420	0.62
nF	7.0	—	—	—	—	—	—	—	—
FA	0.1	1491	0.80	2198	0.82	1809	0.99	5498	0.87
nF	7.0	—	—	—	—	—	—	—	—
AA	0.1	786	1.02	1004	1.29	1020	1.13	2810	1.17
nF	7.0	—	—	—	—	—	—	—	—
HA	0.01	1263	1.03	1466	0.68	852	0.70	3680	0.79
mutation control	—	—	—	—	—	—	—	2584	0.15

¹ number of chromosomes tested. — ² percentage of sex-linked lethals found.

abbreviations:

aF = acid formaldehyde
nF = neutral formaldehyde
FA = formic acid

AA = acetic acid
HA = hydrochloric acid.

All mutation frequencies are significant at the $p < 0.001$ level when compared with the non-treated control.

Table 34. Formic Acid-induced Mutations in *Drosophila Melanogaster*, Larval Feeding Experiments A (450)

treatment	concentration	brood							
		1st		2nd		3rd		4th	
		n ¹	% ²	n	%	n	%	n	%
nF	0.025	910	0.77	717	1.40	737	0.68	2364	0.93
nF	0.05	1266	1.26	1503	0.33	1601	0.31	4370	0.59
aF	0.025	1287	0.79	1213	0.91	1039	0.73	3589	0.78
aF	0.05	754	1.96	802	1.75	901	0.78	2457	1.18
FA	0.1	736	1.15	522	1.34	571	0.88	1879	1.11
AA	0.1	981	0.61	1107	0.60	1157	0.43	3145	0.54
HFA	0.01	757	0.49	663	0.75	800	0.50	2215	0.55
nF	0.025	839	0.95	357	1.17	587	1.19	2283	1.09
aF	0.05	981	1.33	1352	0.96	1571	0.76	3907	0.97
nF	0.025	716	0.93	782	1.28	652	0.77	2150	1.02
aF	0.1	770	0.91	940	1.79	1092	1.10	2702	1.25
nF	0.025	478	1.05	594	1.18	548	0.91	1620	1.05
aF	0.01	416	2.03	514	1.95	447	1.12	1407	1.71
Mutation control	—	—	—	—	—	—	—	2584	0.15

¹ number of chromosomes tested. — ² percentage of sex-linked lethals found.

Abbreviations:

nF = acid formaldehyde

AA = acetic acid

aF = neutral formaldehyde

HFA = hydrochloric acid.

FA = formic acid

All mutation frequencies are significant at the $p < 0.001$ level when compared with the non-treated control.

Table 35. Formic Acid-induced Mutations in *Drosophila melanogaster*, Larval Feeding Experiments B (450)

substance	concentration	pH of the medium	tested chromosomes	mutations found	mutation frequency
nF	0.025%	7.5	398	0	0.00%
nF	0.05%	7.5	366	1	0.27%
FA	0.1%	7.5	544	2	0.38%
Mutation control	—	7.0	2584	4	0.15%

For these experiments a special prepared medium of pH 7 was used (see "Materials and Methods"). Only the 2nd brood was analyzed. Concentration of the glycine-NaOH buffer: 0.1 M. The mutation frequency shows no significant deviation from the (untreated) control (0.15%) in contrast to the results obtained by testing the same substances in an acid medium.

Chemical analysis indicated that the sodium formate administered was completely eliminated, apparently by oxidation, by the 10-12 day of hatching. Survival rates and final embryo weight were essentially the same for both treated and control eggs. No qualitative or quantitative differences in malformations between treated and control embryos were noted. On the other hand, hydrocortisone, included in the study as a positive teratogenic control compound, at a level of 0.025 mg per egg, caused a highly significant increase in malformations in comparison with the controls.

D. Mice

In a study of the correlation between induction of epidermal hyperplasia and tumor development, Frei et al. (142) painted the ears of inbred male Swiss mice (six to ten weeks of age), ten per group, with 8% formic acid in distilled water, twice weekly for periods of 2, 5, 10, 20, and 50 days. At the end of each painting period, all animals were sacrificed, the ears removed, fixed in formalin, and embedded in paraffin. Sections were then cut, stained with hematoxylin and eosin, and examined for increased epidermal cell number (hyperplasia); for variation in thickness of epidermis; and for the presence of an inflammatory exudate. Untreated controls were included in each test. The data were analyzed statistically using the t test. No statistically significant histologic changes resulted.

BIOCHEMICAL ASPECTS

I. Breakdown

Very little information was found in the literature concerning breakdown of formic acid or formates spontaneously or in plants. Formic acid is reported to deteriorate somewhat in normal storage, creating a hazard. Ethyl formate is less stable; it decomposes when dissolved in water.

In sheep, formic acid is rapidly broken down to methane and carbon dioxide by rumen bacteria (482). In man also, part of the formic acid ingested in foods may be dissimilated by microorganisms of the intestinal tract (491).

II. Absorption-Distribution

Formic acid is absorbed from the gastrointestinal tract, through the lungs, through the intact skin, and from the urinary bladder. Ethyl formate is absorbed through the lungs and from the gastrointestinal tract; absorption through the intact skin has not been proven (271,491). Sodium formate is rapidly absorbed from the stomach. Ten minutes after oral administration of 4.44 grams to a human subject, 11.8 mg/100 ml of formic acid was found in the plasma (277). Formic acid, on entering the blood stream, is immediately transformed into sodium formate (277).

Absorbed formic acid is apparently oxidized, partly excreted unchanged in the urine, and partly metabolized in the tissues (491). Retained formate is rapidly utilized in the formation of proteins, lipids, and nucleic acids (370). These in turn are distributed throughout all tissues of the body. Following injection of C^{14} labeled sodium formate, fats from the testes, lungs, spleen, heart, and kidneys had the highest concentration of C^{14} ; depot fats and spinal cord had the lowest. Among the tissue proteins, the greatest amounts of C^{14} were found in the stomach, spleen, kidneys, testes, and liver (434).

III. Metabolism and Excretion

A. Oxidation

The oxidation of formic acid in the organism is affected by several factors such as species, concentration, and certain vitamin deficiencies (105,491). Small doses administered to experimental animals usually are oxidized almost completely whereas 50% or more of larger amounts may be excreted in the urine.

The mechanism of formate oxidation, in rat liver and jejunum at least, involves a catalase-hydrogen peroxide complex and enzymes responsible for H_2O_2 formation, such as xanthine oxidase, uricase, monoamine oxidase, and D-amino acid oxidase (331). In folic acid deficiency, formic acid is not metabolized properly and an increased amount is excreted in the urine (197, 363). Vitamin B_{12} also plays a role in the oxidation of formate; a deficiency increases formic acid excretion but less so than a folic acid deficiency (320, 447). The liver appears to be the main site of formic acid oxidation. Intestinal mucosa, spleen, kidneys, lungs, and blood also oxidize formate. Erythrocytes (frog, chicken, dog, human) also oxidize formic acid aerobically to carbon dioxide by means of the catalase-peroxide complex. In the erythrocyte, peroxide is supplied from a non-enzyme oxidation of certain sulfhydryl compounds like glutathione, homocysteine, and cysteine (371).

The biological half-life of formic acid in various species, according to Malorny (275, 277), are given in Table 36.

Table 36. Biological Half-life of Formic Acid in Various Species (275, 277).

Species	Biological Half-life (minutes)
Rats	12
Guinea-pig	22
Rabbit	32
Cat	67
Dog	77
Man	45 - 46

B. Excretion

1. Rats

Sperling (434) injected male Osborne-Mendel rats, 12-18 months of age, weighing 400-537 grams, intraperitoneally with a single dose of C¹⁴-labeled sodium formate (0.07 - 0.10 mc) in a study of the comparative distribution and excretion of C¹⁴-labeled carbonate and formate in the species. Approximately 80% of the injected dose was excreted over a period of 24 hours.

2. Rabbits

a. Lund (272) injected two full-grown male rabbits, weighing 3150 and 3300 grams, subcutaneously with single doses of sodium formate, 317 and 303 mg/kg BW, respectively. He determined the blood level, rate of formic acid excretion, and toxic effects over a period of 48 hours. One animal was maintained on an acid-forming diet (oats and water); the other on an alkalizing diet (turnips). The results are presented in Tables 37 and 38.

The author concluded from the data obtained that the rabbit is able to oxidize formic acid almost quantitatively and that alkalotic animals excrete larger amounts than acidotic animals.

b. Bastrup (041) injected one rabbit, 2700 grams BW, (strain, age, and sex not given), intramuscularly with formic acid, three doses of 550 mg each, at 24-hour intervals and determined the amount of formic acid excreted in the urine at regular intervals, as well as toxic effects, over a period of 4-5 days.

The animal tolerated 1650 mg of formic acid in divided doses over a period of two days without any apparent ill effects. Approximately 12% of the acid injected was excreted in the urine. Five days following the first injection, the urine level had returned to normal.

c. von Oettingen (491) reported results of an early study in rabbits where the urinary excretion was 14-19% following doses of one g/kg BW.

3. Dog

a. Lund (271) injected a full-grown male dog, 10.1 kg BW, (strain, sex and age not given) subcutaneously with 198 mg/kg BW of sodium formate and determined the blood level and amount of formic acid excreted over a 27-hour period. The animal was observed also for toxic effects. Results obtained are presented in Table 39. The data obtained revealed that 58% of the sodium formate injected was oxidized and 42% was excreted unchanged in the urine.

Table 37. Rabbit No. 2360, Weight 3150 g. Oct. 16, 10 o'clock; subcutaneous Injection of 10 ml 10% Solution of Sodium Formate (\sim 676 mg formic acid). Diet during the Experimental Period: Turnips (272)

Hours aft. Injection	Diuresis ml	Formic acid excret.		Formic acid in blood mg/100 ml	pH
		mg/100 ml	mg total		
3	—	—	—	20	—
4	100	36.8	36.8	16	8.24
10	115	5.5	6.3	10	8.54
23	250	3.0	7.5	—	8.43
32	190	1.4	2.6	—	8.15
48	65	1.0	0.7	—	8.41

Total excretion: 53.9 mg formic acid.

Table 38. Rabbit No. 1. Weight 3300 g. Oct. 19, 11 o'clock; Subcutaneous Injection of 10 ml 10% Solution of Sodium Formate (\sim 676 mg formic acid). Diet during the Experimental Period: Oats (272)

Hours aft. injection	Diuresis ml	Formic acid excreted		Formic acid in blood mg/100 ml	pH
		mg/100 ml	mg total		
—	60	3.2	2.2	—	7.58
—	15	—	—	—	6.21
0	40	—	—	—	5.67
1	—	—	—	48	—
3	10	81.3	8.1	21	5.48
9	10	17.9	1.8	< 10	5.07
26	50	3.8	1.9	—	5.46
46	55	2.1	1.1	—	5.02

Total excretion: 12.9 mg formic acid.

In a second experiment, a single dose of 100 mg/kg BW of sodium formate injected into the washed, empty bladder through a catheter tube, was re-absorbed over a 5-hour period. Preliminary experiments revealed that when sodium formate was given orally, 42% to 56% was excreted unchanged in the urine.

Table 39. The Excretion of Formic Acid after Subcutaneous Administration of Formic Acid

Dog. "H" Weight 10.1 kg. Oct. 30, 10 o'clock: Subcutaneous Injection of 40 ml of a 5% solution of sodium formate (\sim 1.35 g formic acid). (270)

Hours after injection	Diuresis ml	Formic acid in urine		Formic acid in blood mg 100 ml
		mg/100 ml	mg total	
0	2	--	--	--
1	--	--	--	14.2
3.7	48	720	345	15.5
9.5	46	466	215	8.1
23	148	7.5	11	--
27	22	0.0	0.0	--

Total excretion: 0.571 g formic acid.

b. von Oettingen (491) reviewed early studies where following oral administration to dogs of 20, 5, and 1 gram of formic acid, 26%, 65%, and 8% to 9%, respectively, was excreted unchanged in the urine.

4. Man

von Oettingen (491) reports that of 20 grams of sodium formate taken per os by human subjects over a period of two days, 18% was excreted unchanged in the urine. Following an intravenous injection of four grams, 25 to 50% was excreted.

Formic acid is also a normal constituent of human urine of individuals on a mixed diet. Amounts ranging from 13 to 120 mg excreted per day have been reported (156,277).

C. Intermediary Metabolism

The participation of the formate ion in intermediary metabolism is well established (011,156). According to Rappoport (370), "formate is a precursor of serine, methionine, cysteine, purines, and ultimately becomes incorporated, in the form of these precursors, into proteins and nucleic acids". In newborn rats, formate is rapidly metabolized in the formation of proteins, lipids, and nucleic acids. Uptake of C^{14} -formate was found to be high early in life but decreased with age. More formate was incorporated into RNA than into DNA. Black et al.(056) found that there was actually a "preferential utilization" of formate for the synthesis of milk proteins. Fifty percent of C^{14} -labeled formate transferred to milk products was found in the casein.

Annison (011) points out that formate is incorporated into carbohydrates also via glucose by virtue of its being a precursor of the glucogenic amino acid serine involving activation by reaction with tetrahydrofolic acid.

Sperling (434) found that in rats injected intraperitoneally with C^{14} -labeled sodium formate, the retained C^{14} was distributed throughout all tissues of the body.

According to Black et al.(056), although only very small amounts of formate are ingested with foods, many compounds contribute to the C_1 pool for formation of endogenous formate. Serine, glycine, histidine, methionine, choline, and acetone were found to contribute carbon for the synthesis of formate in the rat. Also, other amino acids that give rise to the formation of ketone bodies, such as tyrosine, phenylalanine, or leucine, are potential sources of formate carbon through acetone formation. The liver appears to be an important site of formate metabolism.

Gley et al.(156) also point out that endogenous formic acid is formed through the metabolism of certain amino acids, especially histidine. Glutamic acid, formic acid, and ammonia are produced as these are metabolized in the liver.

Wheeler et al.(506) have outlined the important steps in formate metabolism in their report on the metabolism of normal rat liver and hepatomas (see Fig. 9)




Figure 9. A Simplified Flowsheet of the Metabolism of Formate, Adenine, and Hypoxanthine (506)

Key

Ad	- Adenine	IMP	- Inosinic acid
AMP	- Adenylic acid	HxR	- Inosine
ADP	- Adenosine diphosphate	Hx	- Hypoxanthine
ATP	- Adenosine triphosphate	Xa	- Xanthine
GTP	- Guanosine triphosphate	UA	- Uric acid
GDP	- Guanosine diphosphate	All.	- Allantoin
GMP	- Guanosine monophosphate	XaR	- Xanthosine

The metabolism of C_1 compounds, formate and formaldehyde, in a wide variety of plants and animals, is controlled by folic acid (197). In the folic acid-deficient animal, formic acid is metabolized poorly and large amounts are excreted unchanged in the urine (185,363). On the other hand, in cells with an elevated level of this growth factor, there is an increased incorporation of formate into nucleic acids (197).

Vitamin E deficiency in the monkey also causes a derangement in formate metabolism (105). Severe deficiencies lead to a nutritional muscular dystrophy with anemia and leucocytosis. In this condition there is an increased incorporation of formate into the DNA of skeletal muscle and bone marrow. Vitamin E deficiency does not affect its incorporation into proteins or into nucleic acids to tissues other than the bone marrow or skeletal muscle.

Ascorbic acid deficiency also affects formate metabolism by depressing utilization of formate in the synthesis of serine by 78% and in that of methionine by 85% (320).

According to von Oettingen (491) "the fate of ethyl formate in the organism and its excretion have evidently not been studied."

IV. Effects on Enzymes and Other Biochemical Parameters

Formic acid has an inhibitory effect on certain enzymes in vivo as well as in vitro. Lysozyme, ribonuclease, and trypsin are inactivated (423). Amounts as low as 6 mg/kg BW in the rabbit cause a persistent methemoglobinemia which appears to be due to the action on catalase. On the other hand, gastric secretion and peristalsis are stimulated by formate (491).

von Oettingen (491) has reviewed the literature pertaining to the physiologic effects of formic acid and formates. The nervous system apparently is especially susceptible to the action of formate. In mice, 1 ml of a 2.5% solution of formic acid intrarectally slowed respiration and caused dyspnea, paralysis and death. A dose of 0.46 gm/kg formic acid in the rabbit intravenously depresses the CNS; larger doses caused convulsions. Fatal doses cause clonic, and sometimes tonic convulsions, progressive dyspnea, and paralysis of the medulla. In cats, 400 mg/kg of sodium formate subcutaneously produced somnolence. Larger doses caused staggering and death in about 3 hours. Dogs manifested a staggering gait and weakness lasting 24 hours following subcutaneous administration of 800 mg/kg of sodium formate. Fatal doses intravenously caused clonic convulsions, progressive dyspnea and paralysis of the medulla.

Formic acid at a concentration of 1:1,000,000 has a light stimulating effect on the isolated frog heart, causing a slight increase in amplitude and heart beat. Concentrations as low as 1:10,000, however, quickly and irreversibly arrest heart action (491).

In vivo, formic acid also causes injury to the myocardium that is directly proportional to the dose. The main symptoms are "moderate primary increase and subsequent progressive decrease of the systole, slowing of the heart beat, contraction irregularities, arrhythmias and diastole arrest" (491). Formic acid is considered by some authorities to be seven times more toxic on heart action than formaldehyde, and 187 times more toxic than methanol (491).

With regard to the vascular system, formic acid intravenously causes immediate vasoconstriction. Sodium formate, on the other hand, causes vasodilation in the liver, brain, and kidney but vasoconstriction in the limbs. In the frog, trace amounts of formic acid (1:1,000,000) cause slight vasodilation; dilutions as high as 1:500,000 produce vasoconstriction; and a concentration of 1:1000 causes complete blocking of the circulation. There is a rise in the blood pressure due to intravenous formic acid during vascular constriction; larger doses, however, have a hypotensive effect (491).

Formic acid stimulates gastric secretion and peristalsis and enhances absorption of protein from the intestinal tract. It also exerts a diuretic effect but is less active in this respect than theobromine. Single large doses, or repeated small doses of formate cause kidney injury. Rabbits are the most susceptible of the experimental animals. Diet appears to play a role since animals on an acid diet are more susceptible than those on green foods (491).

Formic acid is reported to possess strong germicidal properties with dilutions as high as 1:50,000 to 1:100,000 being effective against some microorganisms. Concentrations of 60 mg/liter prevent the growth of Bacillus anthracis and B. pyocyaneus. Yeasts, however, are somewhat more resistant, concentrations of 0.2% being required to prevent growth and fermentation. Germicidal action is believed to be due to the un-ionized molecule (491). Formic acid has been used in certain countries as a food preservative (137). Berard et al. (1951) report that a formic acid (20%) colloid of methylolmelamine (17%) prevents microbial deterioration of cotton at a level of 12% resin add-on.

Methyl, ethyl, and isopropyl formates are effective against various life cycle stages of the dried fruit beetle, raisin moth, and other insects that infest dried fruits, nuts, cereals, and tobacco. These compounds are used in this area of the food industry as fumigants (018,139,485,491).

Biehl (054) states "ants have been used in therapy since antiquity". Alcoholic extracts and boiled mixtures of whole ants containing formic acid as the active principle have been used as rubbing ointments, in baths, and also in internal medicine. Depression, melancholia, and nervous weakness (vegetative dystonia) were treated as long as 1000 years ago in this way. Later, these natural concoctions were replaced with synthetic formic acid preparations. Doses of sodium formate as high as three to four grams per day have been used (156). Today, the use of formic acid in medicine is limited mainly to primarily allergic diseases, particularly asthma (non-cardiac type), eczema, psoriasis, urticaria, hay fever, exudative diathesis, rheumatic disease, many types of migraine, pregnancy vomiting, mucitis of the gastrointestinal tract, and chronic kidney inflammation. Unfortunately, treatment for a long time is usually required. Some patients develop hypersensitivity to the compound. Conditions in which large doses of formic acid should not be used are hypertonia, coronary sclerosis, and angina pectoris (054).

Eisenstadter (121) reported good results in the treatment of inflammatory and tumor diseases of the female genital organs with formic acid administered intravenously in doses of 1 ml of a 1:1,000,000 solution every 10 - 14 days for periods up to 10 weeks. In a few acute and subacute cases, doses ten times stronger were safely used.

Meiselas (292) stated that formic acid alone, or in combination with sillicic acid and cobra venom (Nyloxin) had no better effect than an inert control substance in alleviating pain in a double-blind study with twenty-nine osteoarthritis patients.

Formic acid and sodium formate have been used as strong caustic astringents and counter-irritants (438).

V. Drug Interaction

Folic acid antagonist drugs, such as methotrexate, inhibit the metabolism of formic acid resulting in increased amounts being excreted unchanged in the urine (275).

VI. Consumer Exposure Information

Formic acid occurs naturally in a number of food plants, fruits, fruit juices, and wines as well as in milk, cheese, meat, certain vegetables, and some mineral waters (056,146,275,289,303).

In the food industry, formic acid finds use as a flavor adjunct in ice cream, ices, candy, baked goods and some other foods. Concentrations up to 18 ppm are used in certain types of candy (139). Usage levels for a number of other food categories are shown in Table 40. Possible daily intake values for various age groups are shown in Table 41. The amount of formic acid imported has declined steadily from more than three million pounds in 1965 to less than 1/4 million pounds in 1972 (See Table 42) (469). Annual poundage data are given in Table 43.

Formic acid is permitted as a preservative for silage in an amount not to exceed 2.25% of the silage on a dry-weight basis or 0.45% when direct cut. The silage should not be fed to live-stock until four weeks after treatment (020).

Ethyl formate occurs naturally in certain fruits, fruit juices, wines, rum, and some other distilled alcoholic beverages. Oranges, apples, pears, honey, orange juice, and coffee extract are typical fruits and beverages where this compound is found (146).

In the food industry, ethyl formate is used as a component of many flavorings, such as, blueberry, raspberry, strawberry, butter, butterscotch, apple, apricot, banana, cherry, grape, peach, plum, pineapple, tutti fruiti, brandy, rum, arrack, sherry, and whiskey (139). It is used in chewing gum, candy, ices, ice cream, baked goods, and a number of other foods (see Table 40). Possible daily intakes range from 0 to 83.7 mg. (Table 41). Total 1970 poundage reported to NAS and FEMA was 29,077 pounds (see Table 43). Ethyl formate is used also as a bulk and package fumigant (fungicide and larvacide) for certain dried fruits, nuts, dry cereals, and tobacco (018,139,491). With raisins and dried Zante currants, the maximum concentration permitted in the final product is 250 ppm (018).

Formic acid and sodium formate are included in the category of substances possibly migrating to foods from paper and paper board food containers (019).

Acceptable daily intake levels (ADI) for formic acid and ethyl formate are 0-5 mg/kg BW (conditional), for each, calculated as total formic acid from all food additive sources (515,516). Maximum allowable concentrations per 8-hour workshift are 9 mg per cubic meter of air for formic acid and 300 mg per cubic meter of air for ethyl formate (016).

Table 40. Comprehensive GRAS Survey Usage Levels (138)

Substance Name	Food Category	Usual Use WTD Mean%	Max. Use WTD Mean, %
Formic Acid *	Baked Goods (R)	1.912903	5.745161
	Frozen Dairy (R)	1.006897	4.331034
	Soft Candy (R)	1.903448	4.968966
	Gelatin pud (R)	0.481481	3.402778
	Bev Typ I(R)	0.826667	0.885593
	Bev Typ II (R)	1.000000	3.000000
	Hard Candy (R)	6.000000	6.000000
	Misc. Unclass.	0.900000	2.733333
Ethyl Formate **	Baked Goods (R)	0.02752	0.02966
	Frozen Dairy (R)	0.00923	0.01020
	Meat Prods (R)	0.00025	0.00100
	Soft Candy (R)	0.01745	0.01838
	Gelatin Pud (R)	0.00923	0.01495
	Bev Typ I (R)	0.00478	0.00550
	Bev Typ II(R)	0.00019	0.00038
	Hard Candy (R)	0.01158	0.01158
	Chewing Gum (R)	0.00915	0.01479
	Misc. unclas.	0.00887	0.05738

* From: Table 4, Usage Levels Reported for FEMA Questionnaire, Substances Not in NAS Appendix (Group III) Regular Foods Only

** From: Table 2, Usage Levels Reported for NAS Appendix A Substances (Group I) Used in Regular Foods (R)

Table 41. Possible Daily Intakes of NAS Appendix A Substances (Groups I & II)
Per Food Category and Total Dietary, Based on Food Consumption by
Total Sample (138).

Substance name	Food category	No. Firms	Age Level	Possible Ave.	Daily Intakes, mg.	
					High A	High B
Formic Acid	Baked goods		0-5 mo	0.006504	0.008608	0.019534
			6-11 mo	0.048588	0.099088	0.145927
			12-23 mo.	0.104253	0.171779	0.313111
			2-65+yr.	0.262450	0.389850	0.788236
	Frozen dairy		0-5 mo.	0.001007	0.004128	0.004331
			6-11 mo.	0.009566	0.026582	0.041145
			12-23 mo.	0.014499	0.034033	0.062367
			2-65+yr.	0.025777	0.062126	0.110874
	Soft candy		0-5 mo.	0.000381	0.003801	0.000994
			6-11 mo.	0.004188	0.012943	0.019320
			12-23 mo.	0.006662	0.017702	0.017391
			2-65+yr.	0.011040	0.033501	0.028820
	Gelatin pud		0-5 mo.	0.000963	0.001300	0.006806
			6-11 mo.	0.006163	0.018681	0.043556
			12-23 mo.	0.006644	0.016178	0.046958
			2-65+yr.	0.009822	0.025278	0.069417
	Bev Typ I		0-5 mo.	0.001984	0.002976	0.002125
			6-11 mo.	0.018765	0.064232	0.020103
			12-23 mo.	0.048005	0.134333	0.047999
			2-65+yr.	0.085973	0.229565	0.092102
	Bev. Typ II		0-5 mo.	0.000000	0.000000	0.000000
			6-11 mo.	0.000100
			12-23 mo.	0.000200
			2-65+yr.	0.032500	0.094400	0.097500
	Hard candy		0-5 mo.	0.000000	0.000000	0.000000
			6-11 mo.	0.000600	0.001800	0.000600
			12-23 mo.	0.001800	0.005400	0.001800
			2-65+yr.	0.003600	0.010200	0.003600
	All categories		0-5 mo.	0.010838	0.020819	0.033789
			6-11 mo.	0.087869	0.223427	0.262262
			12-23 mo.	0.178664	0.379625	0.489627
			2-65+yr.	0.431162	0.844910	1.190549

Table 41 (Cont'd)

Substance name	Food category	No. Firms	Age Level	Possible Ave.	Daily Intakes, mg. High A High B	
Ethyl formate	Baked goods	20	0-5 mo.	0.935680	1.238400	1.008440
			6-11 mo.	6.990080	14.255360	7.533640
			12-23 mo.	14.998400	24.712960	16.164700
			2-65+yr.	37.757440	56.085760	40.693520
	Frozen dairy	18	0-5 mo.	0.092300	0.378430	0.102000
			6-11 mo.	0.876850	2.436720	0.969000
			12-23 mo.	1.329120	3.119740	1.468800
			2-65+yr.	2.362880	5.694910	2.611200
	Meat prods		0-5 mo.	0.002750	0.007250	0.011000
			6-11 mo.	0.051750	0.139500	0.207000
			12-23 mo.	0.755000	0.129750	0.302000
			2-65+yr.	0.196000	0.325250	0.784000
	Soft candy		0-5 mo.	0.034900	0.349000	0.036760
			6-11 mo.	0.383900	1.186600	0.404360
			12-23 mo.	0.610750	1.622850	0.643300
			2-65+yr.	1.014000	3.071200	1.066040
	Gelatin pud.		0-5 mo.	0.184600	0.249210	0.299000
			6-11 mo.	1.181440	3.581240	1.913600
			12-23 mo.	1.273740	3.101280	2.063100
			2-65+yr.	1.882920	4.845750	3.049800
	Bev Typ. I		0-5 mo.	0.114720	0.172080	0.132000
			6-11 mo.	1.085060	3.714060	1.248500
			12-23 mo.	2.590760	7.767500	2.981000
			2-65+yr.	4.971200	13.274060	5.720000
	Bev Typ.II		0-5 mo.	0.000000	0.000000	0.000000
			6-11 mo.	0.000190
			12-23 mo.	0.000380
			2-65+yr.	0.179360	0.123500
	Hard candy		0-5 mo.	0.000000	0.000000	0.000000
			6-11 mo.	0.011580	0.034740	0.011580
			12-23 mo.	0.034740	0.104220	0.034740
			2-65+yr.	0.069480	0.196860	0.069480
	Chewing gum		0-5 mo.			
			6-11 mo.	0.009150	0.009150	0.014790
			12-23 mo.	0.009150	0.027450	0.014790
			2-65+yr.	0.018300	0.036600	0.029580
	All categories		0-5 mo.	1.364950	2.394370	1.589200
			6-11 mo.	10.589810	25.357560	12.302470
			12-23 mo.	20.922160	40.586130	23.672430
			2-65+yr.	48.332070	83.709750	54.147120

Table 42. U. S. Imports of Formic Acid (469)

Year	Net Quantity (lbs)
1965	3,219,943
1966	3,306,947
1967	
1968	2,011,831
1969	272,808
1970	407,199
1971	407,002
1972	189,251

Table 43. Annual Poundage Data for NAS Appendix A Substances (Groups I & II) (138)

Substance Survey No.	# Reptd. to NAS 1960/1970	Poundage Reptd. NAS (Match. Repts. Both yrs)		Total 1970 Poundage Reptd. NAS	# Repts. to FEMA	Poundage Reptd. FEMA 1970 only	Total 197 Poundage Reported NAS+FEMA
		1960	1970				
Ethyl formate NAS 0079 FEMA 2434	*/*	33	33	33	56	29,044	29,077
Formic acid FEMA 24870	5					128	128

- 1 Adler, A. 1927
Über den einfluss von sauren auf das
wasserbindungsvermögen von serum
Kolloid-Z. 43:313-328
- 2 Aebi, H., and F. Portwich. 1959
Formate oxidation and catalase activity in rats
with Walker sarcoma
Helv. Physiol. Pharmacol. Acta 17:189-201
- 3 Al-Shabibi, M.M.A., E.H. Langner, J. Tobias, and
S.L. Tuckey. 1964
Effect of added fatty acids on the flavor of milk
J. Dairy Sci. 47(3):295-296
- 4 Alha, Antti R., Jyrki Raekallio, and Anna L.
Mukula. 1958
Detection of methanol poisoning with special
consideration of the estimation of formic acid
in solid viscera, blood, and urine. Investigation
of 11 fatal cases
Ann. Med. Exp. Biol. Fenn. (Helsinki) 36:444-451
- 5 Allen, G.R., and M.J. Saxby. 1968
Gas chromatography of isomeric fatty acid methyl
esters
J. Chromatogr. 37(2):312-314
- 6 Allen, I.D., and John K. Haken. 1970
Gas chromatography of homologous esters. III.
Influence of stationary phase polarity on
retention of aliphatic esters
J. Chromatogr. 49(3):409-418
- * 7 Andur, M.O. 1960
Response of guinea pigs to inhalation of
formaldehyde and formic acid alone and with a
sodium chloride aerosol
Int. J. Air Pollut. 3:201-220
- 8 Andur, M.O. 1964
The response of guinea pigs to inhalation of
formaldehyde and formic acid
Am. Ind. Hyg. Assoc. J. 25:564-568
- * 9 American Conference of Government Industrial
Hygienists. 1955
Threshold limit values for 1955
Arch. Ind. Health 11(6):521-524
- 10 Andersson, Bengt. 1934
Die co-zy-mase als co-enzym bei enzymatischen
dehydrierungen
Hoppe-Seyler's Z. Physiol. Chem. 225(2/3):57-68
- * 11 Annison, E.F. and R.R. White. 1962
Formate metabolism in sheep
Biochem. J. 84(3):552-557
- 12 Annison, E.F. 1954
Studies on the volatile fatty acids of sheep
blood with special reference to formic acid
Biochem. J. 58:670-680
- 13 Anon. 1961
Food additives. Methyl formate
Fed. Regist. 26:8667
- 14 Anon. 1962
Food additives. Ethyl formate
Fed. Regist. 27:8308
- * 15 Anon. 1961
Standard Methods of Chemical Analysis
D. Van Nostrand Co., Inc, Princeton, N.J.
2:588-590
- * 16 Anon. 1971
Occupation safety and health standards. 1910.93
Air contaminants
Fed. Regist. 36(157):15101-15104
- 17 Anon. 1971
Modern analytical methods
Allg. Fischwirtschaftsztg. 23(9):24
- * 18 Anon. 1973
Food and drugs: ethyl formate
Code of Federal Regulations, 21 CFR 121. 1096
- * 19 Anon. 1973
Food and drugs: substances that are generally
recognized as safe
Code of Federal Regulations, 21 CFR 121.101
- * 20 Anon. 1973
Food and drugs: formic acid
Code of Federal Regulations 21 CFR 121.219
- 21 Aoki, A., and S. Aihara. 1970
Quality of soy sauce during storage in bottles
Seasoning Sci. (Chomi Kagaku) 17(6):201-214
- 22 Arbogast, Marc, Alfred Charles Maillard, and
Edmond Urien. 1965
The evolution of volatile constituents of beer in
the course of principal fermentation and during
aging (French)
Brasserie 20(219):4-7
- 23 Arnstein, H.R. 1960
The effect of vitamin B12 on metabolism of
formate and certain formate precursors in the rat
Biochem. J. 74:616-623
- 24 Arnstein, H.R.V. 1958
Vitamin B12 and the metabolism of formate and
histidine by the rat
Biochim. Biophys. Acta 29:652-653
- 25 Arnstein, H.R.V. 1959
Changes in the metabolism of DL-serine-3-C-14 and
formate-C-14 in the vitamin B12-deficient rat
Biochem. J. 73:23P
- 26 Asao, Yasuo, and Tamotsu Yokotsuka. 1961
Flavor substances in soy sauce. XVIII. Isolation
of C7H12O5 compound in soy sauce, and its
structure
Nippon Nogei Kagaku Kaishi 35:831-837
- * 27 Association of Official Agricultural Chemists.
1960
Methods of Analysis of the AOAC, 9th ed.
AOAC, Washington, D.C. pp. 171-172; 236-238
- * 28 Association of Official Agricultural Chemists.
1965
Methods of Analysis of the AOAC, 10th ed.
AOAC, Washington, D.C. pp. 205-206; 274-279
- 29 Auclair, J.F., and A. Portmann. 1957
Formic acid as a growth stimulant for
Lactobacillus lactis in autoclaved milk
Nature 179:782-783
- 30 Bacci, Vanna, and Giovanni Mannelli. 1967
Minor components of aqueous alcoholic liquids.
IV. Volatile acids in Italian wines
Atti Congr. Qual., 6th 1967:433-442
- 31 Bach, D. 1932
Desinfektionsversuche
Bull. Sci. Pharmacol. 39(34):499-504
- 32 Bach, D. 1932
Untersuchungen über desinfektionsmittel. II.
Eignung der monobasischen fettsauren
Bull. Sci. Pharmacol. 39(34):425-434
- 33 Bach, D., and J. Lambert. 1937
Die wirkung einiger antiseptika auf die
dehydrogenasen des Saphylococcus aureus. Glucose
aktivierende systeme. Ameisensäure und
eine kleine zahl anderer substrate
Compt. Rend. Seances Soc. Biol. Fil. Assoc.
126:300-302
- 34 Bailenger, J. 1962
A method of separation and identification of
organic acids in feces
Bull. Soc. Pharm. Bordeaux 101(3):137-142

- 35 Baldi, G. 1953
Occupational disease due to formic aldehyde and formic acid
Med. Lavoro 44:485-496
- 36 Ballotta, Francesco. 1936
Chemico-toxicological investigations of formic acid
Boll. Chim.-Farm. 75:577-580
- 37 Barnett, A.J.G., and G.A. Tawab. 1957
The lactic acid and volatile fatty acid contents of five locally made type cheeses
J. Dairy Res. 24(2):225-234
- 38 Bartos, J. 1960
On the utilization of S-(2,5-dihydroxyphenyl)-isothiourea for the identification of formic acid
Ann. Pharm. Fr. 18:907
- 39 Bastianutti, Ione. 1962
Formic acid, oxalic acid, and 8-quinolinol as antiseptics in wines; paper chromatographic analysis
Boll. Lab. Chim. Prov. 13:16-22
- 40 Bastianutti, Jone. 1967
Formic acid in raspberry. Quantitative determination
Ind. Aliment. 6(27):55-58
- * 41 Bastrup, J.T. 1947
On the excretion of formic acid in experimental poisoning with methyl alcohol
Acta Pharmacol. Toxicol. 3:312-322
- 42 Bates, Robert C., and Ronald E. Hurlbert. 1970
The effect of acetate on *Euglena gracilis* var. *bacillaris* as a function of environmental conditions
J. Protozool. 17(1):134-138
- 43 Baum, P., and H. Lamm. 1963
Nonspecific detection of preservatives in various foods
Ernaehrungsforschung 8(3):355-361
- 44 Beacham, L.M. 1946
Decomposition in canned oysters and clams
J. Assoc. Offic. Agric. Chem. 29:89-99
- 45 Becker, Erich. 1954
Paper chromatographic detection of water-soluble organic acids in foods
Z. Lebensm.-Unters. -Forsch. 98:249-257
- 46 Belehradek, J., and P. Schwarz. 1928
L'action toxique des acides de la serie formique en relation avec l'adaptation de l'organisme.
(Toxic action of acids of the formic series in relation to adaptation of the organism)
Bull. Soc. Chim. Biol. 10:909-919
- 47 Bellucci, G., and B. Griqatti. 1970
Separation and determination of some organic acids formed during the spontaneous aerobic fermentation of tomatoes
Ind. Conserve 45(1):11-14
- 48 Benedict, E.M., and G.A. Harrop. 1922
Estimation of formic acid in urine
J. Biol. Chem. 54:443-450
- 49 Benes, L., and V. Drasil. 1962
Incorporation of formate-C-14 and orthophosphate-P-3 in bone marrow cells in vitro. I. Effect of cytosine deoxyriboside 5'-phosphate
Folia Biol. 8(1):40-47
- 50 Bennett, M.A., and G. Toennies. 1942
Nutritional assay of casein modified by action of hydrogen peroxide and formic acid
J. Biol. Chem. 145:671-677
- 51 Berard, W.N., E.K. Leonard and W.A. Reeves. 1961
Cotton made resistant to microbiological deterioration using formic acid colloid of methylolmelamine
Dev. Ind. Microbiol. 2:79-91
- 52 Bergmann, Felix, and Ruth Segal. 1956
The separation and determination of microquantities of lower aliphatic acids, including fluoroacetic acid
Biochem. J. 62(4):542-546
- 53 Bernhauer, Konrad, and Franz Slanina. 1934
Chemismus und enzymchemie der saurebildungs- und saureumwandlungsvorgange bei schimmelpilzen. II. Mitt. Die oxalsaeurebildung aus Ameisensaure, Glykolsaeure, Bernsteinsaure und anderen sauren durch *Aspergillus niger*
Biochem. Z. 274:97-111
- * 54 Biehl, H. 1968
Formic acid in therapy (Ger.)
Quart. J. Crude Drug Res. 8(1):1170-1172
- 55 Bischofswerder, Kurt. 1931
Zweideutige chemische reaktionen in der nahrungsmittelchemie
Mineralwasser-Fabrikant 5:731
- * 56 Black, A.L., M. Kleiber, J.R. Luick, and J.J. Kaneko. 1962
Transfer of carbon from formate to amino acids of milk protein in the intact cow
Am. J. Physiol. 203:1067-1070
- 57 Bleyer, B., W. Diemair, and K. Leonard. 1933
Über den einfluss von konservierungsmitteln auf enzymatische vorgänge
Arch. Pharmaz. Ber. Dtsch. Pharmaz. Ges. 271:539-552
- 58 Blondin, George A., William J. Vail, and David Ezra Green. 1969
Mechanism of mitochondrial swelling. II. Pseudoenergized swelling in the presence of alkali metal salts
Arch. Biochem. Biophys. 129(1):158-172
- 59 Bodansky, Meyer. 1928
Lipoidlöslichkeit, permeabilität und hamolytische wirkung von gesättigten fettsauren
J. Biol. Chem. 79:241-255
- 60 Bohm, E. 1962
A simple ground-glass apparatus for determination of volatile acids in wines and juices
Deut. Lebensm.-Rundsch. 58:255
- 61 Booth, Roland G. 1960
Preservation of whale meat
U.S. Pat. 2,934,440 issued Apr. 26, 1960
- 62 Bosz, J.E. Quintus. 1919
Über oxalsaeure im zuckerrohr und über rohrwachs
Arch. Suikerind. Nederland. Indie 1920:969-977
- 63 Bottazzi, V., B. Battistotti, and M. Vescovo. 1971
Continuous production of yoghurt cultures and stimulation of *Lactobacillus bulgaricus* by formic acid
Milchwissenschaft 26(4):214-219
- 64 Bruce, William F. 1934
Die spaltung von citronensaure durch *Bacillus aertrycke*
J. Biol. Chem. 107:119-129
- 65 Budilova, E.V. 1961
Action of gamma radia ion on deoxyribonucleic acid (DNA) synthesis in isolated nuclei of the thymus. I. Formate-C-14 and adenine-C-14 incorporation into nitrogen bases of the DNA of isolated nuclei
Radiobiology 1:333-335
- 66 Buenadicha, P. 1964
The essential oil of *Cistus ladaniferus*
Farmacognosia 24(3-4):85-104
- 67 Burmeister, H. 1958
Conservation of fruit juices
Brantweinwirtschaft 80:523-526

- 68 Calvarano, Maria. 1966
Composition of Calabrian geranium essence
Essenze Deriv. Agrum. 36(2):100-117
- 69 Canic, Velimir D., and Nada Perisic-Janjic. 1970
Separation of organic acids by thin-layer
chromatography on starch
Tehnika 25(2):330-332
- 70 Cantoni, C., M.A. Bianchi, P. Rennon, and S.
D'Aubert. 1969
Putrefaction of sausages
Arch. Vet. Ital. 20(4):245-264
- 71 Carangal, Apolinario R., Jr., Sylvia R. Arnaldo,
and Erlinda A. Tucay. 1956
Chemical studies on coffee grown in the
Philippines. I. The nonnitrogenous organic acids
of the leaves, shells, and beans
Philipp. Agric. 40:99-103
- 72 Carroll, R.J., and R.E. Hungate. 1955
Formate dissimilation and methane production in
bovine rumen contents
Arch. Biochem. Biophys. 56(2):525-536
- 73 Castagnou, I.R., and J. Zizine. 1951
Detection of cations in toxicology
Bull. Trav. Soc. Pharm. Bordeaux 89:152-155
- * 74 Castle, M.E. and J.N. Watson. 1970
Silage and milk products, a comparison between
wilted and unwilted grass silage made with and
without formic acid
J. Brit. Grassl. Soc. 25(4):278-284
- 75 Castle, M.E., and J.N. Watson. 1970
Silage and milk production, a comparison between
grass silages made with and without formic acid
J. Brit. Grassl. Soc. 25(1):65-70
- 76 Cats, H., and H. Onrust. 1958
Identifying food preservatives by paper
chromatography
Chem. Weekbl. 54:456-459
- 77 Chandra, Suresh. 1969
Microdetermination of formic and oxalic acids in
mixtures with silver (III) and cerium (IV)
solutions
Chim. Anal. 51(11):575-577
- 78 Chang, S.S. 1961
A new technique for the isolation of flavor
components from fats and oils
J. Am. Oil Chem. Soc. 38:669-671
- 79 Chaudhary, S.S., R.E. Kepner, and A.D. Webb. 1964
Identification of some volatile compounds in an
extract of the grape, *Vitis vinifera* var
Sauvignon blanc
Am. J. Enol. Vitic. 15(4):190-193
- * 80 Chemical Rubber Company. 1967
CRC Handbook, 3rd ed.
Chemical Rubber Company, Cleveland, Ohio p. 190
- 81 Chrzasczcz, Tadeusz, and Mikolaj Zakomorny. 1935
Über den mechanismus der umbildung von
ameisensäure zu oxal- und kohlenstaure
durch verschiedene schimmelpilze
Biochem. Z. 279:64-75
- 82 Chung, A.C., F.G. Hueter, and J.C. Shaw. 1959
Oxidative metabolic pattern of normal and ketotic
cow liver incubated with 1-C-14-labeled
aliphatic acids
Proc. Soc. Exp. Biol. and Med. 100(3):476-480
- 83 Claren, Otto Bernd. 1942
Über das schicksal verfütterter ameisensäure im
verdauungstrakt des wiederkauers
Hoppe-Seyler's Z. Physiol. Chem. 276(3):97-107
- Clark, E.P., and Fred Hillig. 1938
The Dyer method for the identification and
determination of volatile fatty acids
J. Assoc. Offic. Agric. Chem. 21:684-688
- 85 Clark, J.B. 1953
The mutagenic action of various chemicals on
micrococcus aureus
Proc. Okla. Acad. Sci. 34:114-118
- 86 Clarke, B.J., F.V. Harold, R.P. Hildebrand, and
A.S. Morieson. 1962
Trace volatile constituents of beer. IV. Volatile
acids
J. Inst. Brewing 68:179-187
- 87 Cohn, Robert. 1921
Ein neuartiger nachweis von Formaldehyd mit
resorcinschwefelsaure
Chem. Ztg. 45:997-998
- * 88 Committee on Specifications. 1972
Food Chemicals Codex
Committee on Food Protection, Nat. Acad. Sci.
Nat. Res. Council, Washington, D.C.
- 89 Corbellini, A., G. Lugaro, I. Lupi, and E. Crespi.
1966
Research on Urogastrone. V. The action of formic
acid on the biologically active fraction
Ital. J. Biochem. 15(3):151-171
- 90 Corrao, Andrea. 1957
Researches on the soluble volatile acids of olive
oil
Olivi Miner., Grassi Saponi, Colori Vernici
34:536-538
- 91 Craddock, V.M. 1971
Methylation of DNA in the intact animal and the
effect of the carcinogens dimethyl-nitrosamine
and ethionine
Biochim. Biophys. Acta 240:376-383
- 92 Crosbie, G.W. 1962
Identification of formic acid as the p-
bromophenacyl ester
Nature (London) 193:579
- 93 Cruz, N.C. Vera, J.R. Lodge, and C.N. Graves.
1967
End products of bovine spermatozoan metabolism as
influenced by incubation diluent and temperature
J. Dairy Sci. 50(4):558-561
- 94 Davies, C.W., and B.D.R. Owen. 1956
Adsorption of organic acid molecules and their
chromatographic separation on ion-exchange resins
J. Chem. Soc. 1956:1681-1685
- 95 Delitsch, Heinrich. 1938
Über mikrokokken im tilsiter kase
Zbl. Bakt. Parasitenk. Infektionsk. Abt. II.
97:389-399
- 96 Delore, P., and J. Jeannin. 1928
Untersuchungen über fettsauren. Verhältnis
zwischen chemischer konstitution
und experimentellen schädigungen
Compt. Rend. Soc. Biol. 98:700-702
- * 97 Demerec, M., G. Bertani and J. Flint. 1951
A survey of chemicals for mutagenic action on *E.*
coli
Am. Nat. 85:119-136
- * 98 Deshusses, J. 1961
Formic acid content of roasted coffee, chicory,
soluble coffee extracts, and coffee substitutes
Mitt. Geb. Lebensm. Hyg. 52:428-430
- 99 Deshusses, J. 1961
Fincke's modified method for the determination of
formic acid
Mitt. Geb. Lebensm. Hyg. 52:422-427
- 100 Dhindsa, K.S. 1969
Time course of incorporation of formate-C-14, a
nucleic acid precursor, in mouse liver cells
Indian J. Exp. Biol. 7(2):84-85

- 101 Diemair, W., and C. Gundermann. 1959
Determination of formic acid in wine
Z. Lebensm.-Unters. -Forsch. 110:261-267
- 102 Diemair, W., and P. Schams. 1960
Gas chromatography in food analysis. I.
Determination of the lower, volatile fatty acids
in foods
Z. Lebensm.-Unters. -Forsch. 112:457-463
- 103 Dimitrova, E., V. Staikov, and G. Zolotovitch.
1955
Pelargonium roseum in Bulgaria (Russian)
Compt. Rend. Acad. Bulg. Sci. 8:33-36
- 104 Dinning, J.S., J.T. Sime, and P.L. Day. 1955
Influence of vitamin E deficiency on metabolism
of sodium formate-C-14 and glycine (aminoacetic
acid)-1-C-14 by rabbit
J. Biol. Chem. 217:205-211
- * 105 Dinning, James S., and Paul L. Day. 1958
Vitamin E deficiency in the monkey. III. The
metabolism of sodium formate-C-14
J. Biol. Chem. 233(1):240-242
- 106 Dinning, James S., Lloyd D. Seager, Lou D. Payne,
and John R. Totter. 1952
The metabolism of betaine and sodium formate by
leukemic mice
Science 116:121-122
- 107 Dodge, Francis D. 1941
Geranium oil
Soap Sanit. Chem. 17(3):33, 49
- 108 Doelle, Horst W. 1969
Gas chromatographic separation and determination
of microquantities of C1-7 branched and straight-
chain saturated fatty acids
J. Chromatogr. 39(4):398-406
- * 109 Dowden, Bobby F., and Harry J. Bennett. 1965
Toxicity of selected chemicals to certain animals
J. Water Pollut. Control Fed. 37(9):1308-1316
- 110 Drawert, F., H.-J. Kuhn, and A. Rapp. 1962
Reaction gas chromatography. III. Gas
chromatographic determination of lower fatty
acids in the stomach of leaf-eating monkeys
(Colobinae)
Z. Physiol. Chem. 329(1-2):84-89
- 111 Drews, E. 1954
Detection of organic acids and preservatives by
means of paper chromatography
Getreide Mehl 3:85-88, 90-91
- 112 Drews, E. 1955
Paper chromatography of bread and baked goods.
II. Determination of volatile acids
Brot Geback 9:45-47
- 113 Drews, E. 1956
The origin of formic acid in the baking process,
and its determination
Brot Geback 10:166-171
- 114 Droller, Hugo. 1932
Zum nachweis geringer mengen von ameisensaure in
blut und gewebe, insbesondere in der haut
Z. Physiol. Chem. 211:57-64
- 115 Du Vigneaud, Vincent, Walter G. Verly, John E.
Wilson, Julian R. Rachele, 1951
One-carbon compounds in the biosynthesis of the
"biologically labile" methyl group
J. Am. Chem. Soc. 73:2782-2785
- 116 Dudkin, M.S., and N.G. Shkantova. 1961
Determination of volatile organic acids in bran
of grain hulls
Zh. Priklad. Khim. 34:1337-1342
- 7 Eds, F. 1924
Quantitative verfahren zur bestimmung der
ameisensaure im blut und harn
J. Lab. Clin. Med. 10:59-66
- 118 Ege, Rich. 1924
Untersuchungen uber die durchdringungsfahigkeit
der anionen bei beruhrung mit der membran der
roten korperchen
Compt. Rend. Soc. Biol. 91:779-781
- 119 Egorova, L.N., L.L. Lagunov, and M.I. Kryuchkova.
1967
Preserving meat of marine mammals by acids
Tr. Vses. Nauch.-Issled. Inst. Morsk. Ryb. Khoz.
Okeanogr. 63:81-85
- 120 Egorova, L.N., V.I. Treshcheva, and E.M. Sidorova.
1967
The influence of preservatives on the quality of
fish minced meat and flour produced from it
Tr. Vses. Nauch.-Issled. Inst. Morsk. Ryb. Khoz.
Okeanogr. 63:86-93
- * 121 Eisenstadter, D. 1932
Formic acid treatment of female genital
inflammatory diseases.
Zentralbl. Gynak. 56:161-162
- 122 Eisner, M. 1962
Determination of formic acid in fruit sirups
Ind. Obst.-Gemueseerwert. 47:503-504
- 123 El-Gindy, M. Momtaz, and E. Drews. 1957
Paper-chromatographic identification of some
preservatives in baked products
Brot Geback 10:224-228
- 124 Eliassen, K.A., and O.V. Sjaastad. 1968
Effects of formaldehyde and formic acid on
histamine metabolism in sheep
Acta Pharmacol. Toxicol. 26(6):482-490
- 125 Elsdon, S.R. 1946
Application of the silica gel partition
chromatogram to the estimation of volatile fatty
acids
Biochem. J. 40:252-256
- 126 Emmelin, N., et al. 1947
The mechanism of the sting of the common nettle
(Urtica urens)
J. Physiol. 106:440-455
- 127 Erra, Ugo. 1958
Physical and histopathological findings in cases
of poisoning with methyl alcohol, formaldehyde,
and formic acid
Polia Med. 41:366-377
- * 128 Estienne, Jean. 1964
Honey
Fr. Parfums 7(41):383-390
- 129 Parkas, A., L. Parkas, and J. Yudkin. 1934
Uber die spaltung von natriumformiat durch
Bacterium coli in gegenwart von schwerem wasser
Proc. Royal Soc., Lond., Ser. B. 115:373-379
- 130 Feldberg, W. 1940
The action of bee venom, cobra venom, and
lysolecithin on the adrenal medulla
J. Physiol. 99:104-118
- 131 Fenaroli, Giovanni, Franco Poy, and Giuseppe
Maroni. 1965
Natural and synthetic flavors. I. Rum and rum
ether (Italian)
Riv. Ital. Essenze Profumi Pianta Offic. Aromi
Saponi Cosmet. 47(9):484-495
- 132 Filov, V.A. 1961
Retention and transformation of low-molecular-
weight esters in the organism
Gig. Tr. Prof. Zabol. 5(3):14-19
- 133 Fisher, C.D. 1945
Controlling storage insects by fumigation
Food Ind. 17(10):1176-1178
- 134 Flaschentrager, Bonifaz, and Rudolf v.
Wolffersdorff. 1934
Uber den giftstoff des crotonoles. I. Die sauren
des crotonoles
Helv. Chim. Acta 17:1444-1452

- 135 Flossner, O. 1931
Beiträge zur biochemie menschlicher
körperflüssigkeiten. II. Untersuchungen
über ovarialzystenflüssigkeiten
Z. Biol. 91:291
- 136 Flury, P. 1920
Über die bedeutung der Ameisensäure als natürlich
vorkommendes gift (Ger.)
Ber. Dtsch. Pharm. Ges. 29:650-673
- * 137 Food and Agriculture Organization. 1969
Current food additives legislation - Germany
Curr. Food Addit. Legisl. 1969(121):4
- * 138 Food Protection Committee. 1972
Comprehensive GRAS survey
Nat. Acad. Sci., Nat. Res. Council, Washington,
D.C.
- * 139 Food Protection Committee, Food and Nutrition
Board. 1965
Chemicals used in food processing
Nat. Acad. Sci., Nat. Res. Council, Washington,
D.C., Publication 1274
- 140 Fourton, A. 1928
Existiert die antirrhinsäure in der digitalis?
Bull. Sci. Pharmacol. 35:689-691
- * 141 Freese, E.B., J. Gerson, H. Zaber, H.-J. Rhaese
and E. Freese. 1967
Inactivating DNA alterations induced by peroxides
and peroxide-producing agents
Mutat. Res. 4:517-531
- * 142 Frei, J.V. and P. Stephens. 1968
The correlation of promotion of tumor growth and
of induction of hyperplasia in epidermal two-
stage carcinogenesis
Brit. J. Cancer 22:83-92
- 143 Freund, J. 1933
Sodium formate as substitute for sodium chloride
Klin. Wchnschr. 12:226-227
- 144 Friedmann, B., H.I. Nakada, and S. Weinhouse.
1954
Study of oxidation of formic acid in folic acid-
deficient rat (using C14 labeled formate
J. Biol. Chem. 210:413-421
- 145 Fukui, Kiyoshi, Hideo Nagatomi, and Saburo Murata.
1962
Direct determination of low-molecular-
weight fatty acids by gas-liquid chromatography
Bunseki Kagaku 11:432-434
- * 146 Furia, T.E. and N. Bellanca, (Eds.). 1971
CRC Penarali's Handbook of Flavor Ingredients
The Chemical Rubber Company, Cleveland, Ohio, 1971
- 147 Genov, I. 1967
Effect of bactericidal substances on the
dehydrogenase activity of Bacillus
anthracis Mycobacterium tuberculosis at various
temperatures
Tr. Tsent. Nauch.-Issled. Dezinfek. Inst.
18(Pt. 1):140-144
- 148 Gerlsma, S.Y. 1970
Gelatinizing temperatures of starch, as affected
by polyhydric and monohydric alcohols, phenols,
carboxylic acids, and some other additives
Stärke 22(1):3-9
- 149 Gianturco, V.M., A.S. Giammarino, and P. Friedel.
1966
Volatile constituents of coffee
Nature 210(5043):1358
- 150 Gierschner, K. 1966
Determination of formic acid in fruit juices
Int. Fruchtsaft-Union, Ber. Wiss.-Tech. Komm.
7:135-158
- 151 Gierschner, K. 1968
Methods for determining formic acid in fruit
juices
Ind. Obst. Geaueseverwert. 53(12):355-361
- 152 Giraddinov, Zh.K., L.N. Lishtvanova, and M.I.
Goryaev. 1962
Essential oil from Artemisia porrecta var.
coerulea (Russian)
Izv. Akad. Nauk Kaz. SSR, Ser. Khim.
1962(1):112-116
- 153 Gimenez, E. Hernandez, and B. Lafuente Perriols.
1963
Orange juice preservatives
Rev. Agroquim. Tecnol. Aliment. 3(2):136-142
- 154 Gjerstad, Gunnar. 1965
Intermediary metabolism of ergot. XII.
Comparative metabolism of nicotinic,
mevalonic, acetic, citric, and formic acids,
proline, methionine, and tryptophan
Q. J. Crude Drug Res. 5(3):735-738
- * 155 Gleason, M.N., et al. 1969
Clinical Toxicology of Commercial Products, 3rd
ed.
The Williams & Wilkins Company, Baltimore
- * 156 Gley, Pierre, and Jean-Emile Courtois. 1967
Report from the Commission on Nutrition
concerning the use of formic acid and formates
as agents for preservation of food products.
Bull. Acad. Nat. Med. 151:598-601
- 157 Goepfert, John M., and Ruthann Hicks. 1969
Effect of volatile fatty acids on Salmonella
typhimurium
J. Bact. 97(2):956-958
- 158 Gogiya, V.T., and L.N. Ivanova. 1968
Composition of pink geranium essential oil
Prikl. Biokhim. Mikrobiol. 4(6):733-738
- 159 Golber, L.M. 1935
Formic acid, in cancer patients; diagnostic
significance
Vrach. Delo 18:119-124
- 160 Gould, I.A., and R.S. Prantz. 1946
Formic acid content of milk heated to high
temperatures as determined by the distillation
procedure
J. Dairy Sci. 29:27-31
- 161 Gould, I.A., Earl Weaver, and R.S. Prantz. 1946
Lactic acid in dairy products. IV. Effects of
storage of evaporated milk on lactic and formic
acid production, on pH, and on acidity and formal
titrations
J. Dairy Sci. 29:33-40
- * 162 Gray, J.A. 1966
Kinetics of enamel dissolution during formation
of incipient caries-like lesions
Arch. Oral Biol. 11:397-421
- 163 Green, James W. 1949
The relative rate of penetration of the lower
saturated monocarboxylic acids into mammalian
erythrocytes
J. Cell. Comp. Physiol. 33:247-266
- 164 Greig, Margaret Elizabeth, Muriel Platt Munro,
and Kenneth Allan Caldwell Elliott. 1939
Milch- und brenztraubensäurestoffwechsel in
normalen und tumorgewebe. VI. Rinderretina und
kühnembryo
Biochem. J. 33:443-453
- 165 Groll, Hermann, and G. Borger. 1926
Experimentelle untersuchungen zur lehre von der
entzündung. IV. Mitt. Die sauerstoffatmung des
gewebes bei entzündung und reizung
Krankheitsforschung 2:220-262

- 166 Grossfeld, J., and R. Payfer. 1939
Determination of formic acid in foods
Z. Unters. Lebensm. 78:1-30
- * 167 Gruene, A., and Volker Nobbe. 1967
Preservatives and their identification using
prepared plates for thin layer chromatography
Riechst. Aromen. Koerperpflegen. 17(12):501-514
- 168 Grunhut, L. 1921
Nachweis und quantitative bestimmung der
lavulinsaure in lebensmitteln
Z. Unters. Nahrgrs. Genussm. 41:261-280
- 169 Guagnini, Omar A., and Eugenio E. Vonesch. 1959
Hydroxamic acids. IX. Microdetection and
determination of formic acid and inorganic
formates: application to foods
An. Assoc. Quim. Arg. 47:41-51
- 170 Guetlbauer, P. 1969
Chromatographic behavior and chemical structure.
I. Thin-layer chromatography of aliphatic acids
J. Chromatogr. 45(1):104-112
- 171 Gulati, Baldev C. 1969
Essential oils of wild ginger of Kumaon
Perfum. Essent. Oil Rec. 60(78):257-258
- 172 Gustav, J., Martin Stenzel, and Wolfram Stenzel.
1944
The influence of various chemicals and vitamin
deficiencies on the excretion of glucuronic acid
in the rat
Arch. Biochem. 3:325-331
- * 173 Hagan, E.C., W.H. Hansen, O.G. Fitzhugh, P.M.
Jenner, W.I. Jones, Jean M. Taylor, Eleanor L.
Long, A.A. Nelson, and J.R. Brouwer. 1967
Food flavorings and compounds of related
structure. II. Subacute and chronic toxicity
Food Cosmet. Toxicol. 5(2):141-157
- 174 Hailer, E. 1938
The action of disinfectants on human and bovine
tubercle bacilli. IV. Experiments with aromatic
acids
Z. Hyg. Infektionskr. 121:405-431
- 175 Hall, J. Alfred, and C.P. Wilson. 1925
Die fluchtigen bestandteile von
valenciaorangensaft
J. Am. Chem. Soc. 47:2575-2584
- 176 Hamsik, A. 1929
Action on oxyhemlin and chlorhemlin
Z. Physiol. Chem. 183:103-112
- 177 Hanak, A. 1940
In preservation of raspberry juice
Biochem. Z. 221:467-472
- 178 Hanak, A., and K. Kurschner. 1930
New method of testing for formic acid as
preservative in fruit juices
Z. Unters. Lebensm. 60:278-290
- 179 Harper, W.J., and H.E. Randolph. 1965
Relationship of acids to taste and flavor of
buttermilk
Milk Dealer 55(1):65, 68-9
- 180 Harper, W.J., I.A. Gould, and C.L. Hankinson.
1961
Observations on the free volatile acids in milk
J. Dairy Sci. 44:1764-1765
- 181 Henderson, J. Frank, and G.A. LePage. 1959
Purine biosynthesis de novo in mouse tissues and
a mouse tumor
J. Biol. Chem. 234:2364-2368
- 182 Herken, W. 1968
The influence of blood-pH on ionization,
distribution, and toxicity of formic acid
Naunyn Schmiedeberg Arch. Pharm. Exp. Path.
260:142-143
- 183 Herner, Birger. 1944
Effect of benzoic acid and of related compounds
on the ammonia formation in kidney and liver
tissue of the rat
Acta Physiol. Scand. 8(Suppl. 23):99 pp
- 184 Hess, Ludwig. 1935
Die geschmackliche beeinflussung von obstsaften
durch konservierungsmittel
Obst. Gemuse-Verwertg.-Ind. 22:383-385
- * 185 Hiatt, H.H., J.C. Rabinowitz, R. Loch and M.
Goldstein. 1958
Effects of folic acid antagonist therapy on
urinary excretion of formic acid by humans
Proc. Soc. Exp. Biol. Med. 98(1):144-147
- 186 Hillig, Fred. 1942
Effect of neutralization on sour milk in the
manufacture of dried skim (milk)
J. Assoc. Offic. Agric. Chem. 25:253-264
- 187 Hillig, Fred. 1944
Report on (the analysis of) dried and skim milk
(lactic acid)
J. Assoc. Offic. Agric. Chem. 27:235
- 188 Hillig, Fred. 1944
Report on (the determination of) volatile acids
in fish and fish products
J. Assoc. Offic. Agric. Chem. 27:237-240
- 189 Hillig, Fred. 1945
Report on (the determination of) acid in eggs
J. Assoc. Offic. Agric. Chem. 28:217-221
- 190 Hillig, Fred, and L.L. Ramsey. 1945
Determination of lactic and volatile acids in
fruits and fruit products
J. Assoc. Offic. Agric. Chem. 28:525-530
- 191 Hillig, Fred, L.R. Shelton, Jr., and J.H. Loughrey
1961
Chemical indexes of decomposition in ocean perch
J. Assoc. Offic. Agric. Chem. 44:488-493
- 192 Hillig, Fred, L.R. Shelton, Jr., J.H. Loughrey,
and Sammie Bethea. 1961
Chemical indexes of decomposition in pollock and
whiting
J. Assoc. Offic. Agric. Chem. 44:499-507
- 193 Hillig, Fred, L.R. Shelton, Jr., J.H. Loughrey,
and Jerome Fisher. 1958
Chemical indexes of decomposition in cod
J. Assoc. Offic. Agric. Chem. 41:763-776
- 194 Hohl, L.A., and M.A. Joslyn. 1941
Formic acid formation in alcoholic fermentation
Plant Physiol. 16:755-769
- 195 Hollaender, Alexander, W.K. Baker, and E.H.
Anderson. 1951
Effect of oxygen tension and certain chemicals on
the x-ray sensitivity of mutation production and
survival
Cold Spring Harbor Symp. Quant. Biol. 16:315-326
- 196 Hori, Satoshi, and Akira Tanabe. 1956
Quantitative estimation of straight-chain
volatile fatty acids by column
partition chromatography
Okayama Daiyaku Nogakubu Gakujutsu Hokoku 9:47-54
- * 197 Huennekens, F.M., M.J. Osborn, and H.R. Whiteley.
1958
Folic acid coenzymes. Metabolic reactions
involving "active formate" and "active
formaldehyde" are surveyed
Science 128(3316):120-124
- 198 Huitson, John J., and Peter W.J. Watts. 1968
Prevention of mold in animal feeds
Et. Pat. 1,549,021 issued Dec. 6, 1968
- 199 Imai, Nobukatsu. 1930
Experimentelle untersuchungen uber die
verknoecherung im auge
Graefe's Arch. Ophthalmol. 125(2):267-299

- 200 Imai, S., and K. Suzuki. 1969
Components of shoyu mash brewed at different
temperatures in the early stage
Seasoning Sci. 16(5):1-5
- 201 Inoue, Yoshiyuki, and Manjiro Noda. 1950
Paper partition chromatography of aliphatic
carboxylic acids by means of hydroxamic acid
method
J. Agric. Chem. Soc. Jap. 23:368
- 202 Inouye, Shinichi. 1925
Die kennzeichnung der biologischen wirkung von
sauren auf das froschherz
Acta Scholae Med. Kioto 7:227-240
- 203 Iranzo, J. Royo. 1966
Methods of analysis applicable to concentrated
juices and other products from citrus fruits
Conserve Deriv. Agrum. (Palermo) 15(3):109-114
- 204 Ishigaki, Reisaburo, and Ichiro Nagase. 1964
Semichemical soy sauce. III. Quantitative
transition of amino acids in the
semichemical processing of soy sauce
Nippon Jozo Kyokai Zasshi 59(2):161-162
- 205 Ishikawa, Mitsuteru. 1948
Experimental anaphylaxis induced by simple
chemical compounds. I
Jap. Med. J. 1:385-396
- 206 Ishinda, Shuzo. 1962
Organic acids and fertilization membrane in the
sea urchin, *Anthocardia crassispina*
Chiba Daigaku Bunri Gakka Kiyo, Shizen Kagaku
3(4):495-502
- 207 Issakowa, A. 1934
Über die gegenseitigen beziehungen zwischen
höheren pflanzen und mikroorganismen
Iswestija Akad. Nauk S.S.S.R. 1934(7):993-1006
- 208 Itoh, Michio, Yasuharu Nakamura, and Kazuo Shibata
1962
Inhibition of catalase by carboxylic acids
related to dissociation and association of the
apoprotein
Can. J. Biochem. Physiol. 40:1327-1334
- 209 Iwashita, Seishichi. 1959
Rancidity of miso. III. Compositions of the
volatile acids of rancid miso and normal miso
Hakko Kogaku Zasshi 37:264-267
- * 210 Iyer, Meena, T. Richardson, C.H. Amundson, and
R.C. Tripp. 1967
Major free fatty acids in Gouda cheese
J. Dairy Sci. 50(3):385
- 211 Jackson, R.B. 1964
Gas-liquid chromatography of volatile fatty acids
from formic acid to valeric acid. I. Carboxylic
acids as stationary phases
J. Chromatogr. 16(2):306-310
- 212 Jackson, R.B. 1966
Gas-liquid chromatography of volatile fatty acids
from formic to valeric acid. II. The instability
of silicone oil-fatty acid stationary phases
J. Chromatogr. 22(2):251-260
- 213 Jaiswal, P.K., and S. Chandra. 1969
Microdetermination of formic and acetic acids in
their mixture with copper (III) and silver (III)
solutions
Chim. Anal. 51(10):493-495
- 214 Jancso, N., and A. Jancso-Gabor. 1959
Permanent elimination of chemical pain
sensitivity by capsaicin
Arch. Exp. Pathol. Pharmacol. 236:142-145
- 215 Janke, Alexander, and Ferdinand Beran. 1933
Über die mikrobicide wirkung von organischen
sauren und ihren kupfersalzen. Ein beitrag zum
problem des zusammenhangs zwischen chemischer
konstitution und mikrobicide wirkung
Arch. Mikrobiol. 4:64-71
- 216 Jankovic, Aleksandar, and V. Adamovic. 1955
Determination of sulfur dioxide and formic acid
in fruit products
Tehnika 10:1317-1320
- * 217 Jenner, P.M., E.C. Hagan, Jean M. Taylor, E.L.
Cook, and O.G. Fitzhugh. 1964
Food flavorings and compounds of related
structure. I. Acute oral toxicity
Food Cosmet. Toxicol. 2(3):327-343
- 218 Josefsson, L. 1966
Studies with model substances on the mechanism of
the formic acid-induced reversible inactivation
of protein enzymes
Biochim. Biophys. Acta 115:148-159
- 219 Josefsson, Lars. 1962
Changes in the physico-chemical properties of
proteins during the formic acid-
induced transformation
Biochim. Biophys. Acta 59(1):128-136
- 220 Kabrhel, G. 1923
Preservation of fruit juices
Cas. Lek. Cesk. 62:954-955, 982-985, 1008-1010
- 221 Kapeller-Adler, Regine, and Edith Lauda. 1933
Über atherlosliche sauren im harn bei
verschiedener ernahrung
Biochem. Z. 258(5/6):429-448
- * 222 Karunakaran, Rajyasevanirata C.O., and K.K.
Marayana Pillai. 1944
Formic acid and acetic acid as poisons
Indian Med. Gaz. 79:598-601
- 223 Kashima, Tetsu. 1954
Titration of nonaqueous solutions of drugs and
chemicals. III. The determination of sodium
salts of carboxylic acids in glacial acetic acid
Bull. Natl. Hyg. Lab. (Tokyo) 72:145-150
- 224 Kawai, Yukio. 1968
Free fatty acids of *Ascaris lumbricoides* and
Anisakis species larvae. II. Volatile free fatty
acids of *Ascaris lumbricoides*
Kiseichugaku Zasshi 17(6):481-486
- 225 Kawakishi, Shunro, Susumu Hirano, and Hisao Yoshii
1961
Organic acids formed in process of miso
fermentation. I. Organic acids of soybean koji
Nippon Noeikagaku Kaishi 35:347-350
- 226 Keil, Werner. 1941
The question of the harmfulness of formic acid
added as a preservative in the production of
silage
Biedermanns Zentr. B Tierernahr. 13:211-215
- 227 Keilin, D., and E.P. Hartree. 1955
Catalase, peroxidase and metmyoglobin as
catalysts of coupled peroxidatic reactions
Biochem. J. 60(2):310-324
- 228 Kern, J., H.H. Weiser, W.J. Harper, and I.A. Gould
1954
Organic acids formed during the heat
sterilization of milk
J. Dairy Sci. 37:904
- 229 Kienhuis, H., G. Blasse, and J. Matze. 1959
Action of anhydrous formic acid on peptides and
proteins
Nature (London) 184(Suppl 26):2015-2016
- 230 Kini, M.M., D.W. King, Jr., and J.R. Cooper. 1962
Biochemistry of methanol poisoning. V.
Histological and biochemical correlates of
effects of methanol and its metabolites on the
rabbit retina
J. Neurochem. 9:119-124
- 231 Kirsch, W., H. Jantzon, and P. Sieden. 1943
Preservation of fish by acidification with
potatoes or formic acid (Amasil) and its
utilization by swine
Biedermanns Zentr. Abt. B15:209-228

- 232 Klima, D. 1967
Changes in the contents of volatile flavor compounds during the storage of meat and in spoiling of meat
Tehnol. Mesa 8(3):67-68
- 233 Kluge, Hans. 1933
Über den einfluss von konservierungsmitteln auf die wirkung von fermenten
Z. Unters. Lebensm. 66:412-435
- 234 Knoop, F., and H. Jost. 1923
Über milchsäureausscheidung in harn
Z. Physiol. Chem. 130:338-349
- 235 Kobzarenko, V.S. 1939
Chemical preservatives and their determination in food products
Akad. Nauk. U.S.S.R., Inst. Khim. Tekhnol. 1939:113 pp
- 236 Kolesnikova, I.G. 1959
Rol' murav'inoi kisloty v obmene veshchestv bakterii i drugikh organizmov. (The role of formic acid in the metabolism of bacteria and other organisms)
Usp. Sovrem. Biol. 47(1):49-63
- 237 König, J., and E. Kracht. 1929
Natur und bedeutung der fluchtigen stoffe der nahrungsmittel
Z. Unters. Lebensm. 57:377-412
- 238 Kopaczewski, W. 1934
Die gelatinisierung des serums durch organische säuren
Compt. Rend. Hebd. Seances Acad. Sci. 198:1271-1273
- 239 Kovacs, A.S., and P. Denker. 1962
Identification and determination of food preservatives in foods
Ind. Obst.-Gemüseverwert. 47(1):1-3
- 240 Kovacs, Akiva S., Peter Denke, and Hans O. Wolf. 1962
Detection and determination of preservatives in food. II. Determination of food preservatives
Ind. Obst.-Gemüseverwert. 47:547-559
- 241 Kramer, Magdalene, and R. Tarjan. 1962
Effects of preservatives on the utilization of carotene
Int. Z. Vitaminforsch. 32:149-157
- 242 Kreutz, A., and C. Buchner. 1926
Formic acid in table vinegar and essence of vinegar
Z. Unters. Lebensm. 52:295-298
- 243 Krylova, N.N., V.P. Volovinskaya, V.M. Gorbatov, I.N. Lyaskovskaya, K.I. Bazarova, R.T. Khlamova, and G.A. Yakovleva. 1969
Elimination of smoke in sausage production
Tr. Vses. Nauchno-Issled. Inst. Mясnoi Prom-st. 22:17-31
- 244 Kukushkina, G.V., et al. 1966
Inhibition of biosynthesis of protein and nucleic acids by phenols in vivo
Vopr. Med. Khim. 12:452-455
- 245 Kulman, J., and R. Zadina. 1931
Harmful effect
Casop. Lek. Cesk. 70:985-988
- 246 Kulshrestha, D.C., and E.H. Parth. 1970
Inhibition of lactic streptococci and some pathogenic bacteria by certain milk-associated volatile compounds as measured by the disk assay
J. Milk Food Technol. 33(8):305-310
- 247 Kumar, Kamal, V.N. Sarin, and Putcha Venkateswariu. 1969
Infrared spectrum of solid ethyl formate
Proc. Nucl. Phys. Solid State Phys. Symp., 13th 1968(3):230-233
- 248 Kuntze, K., and W. Peyer. 1947
Quick testing and assaying of drugs
Suddeut. Apoth.-Ztg. 87:226-227
- 249 Kuvaev, B.E., and N.S. Imyanitov. 1967
Determination of monobasic and dibasic acids by paper chromatography
Zavod. Lab. 33(11):1382-1383
- 250 Lafon, Madeleine. 1955
Formation of secondary products of alcoholic fermentation
Ann. Inst. Natl. Recherche Agron., Ser. E., Ann. Technol. Agr. 4:169-221, 241-292
- 251 Lagunov, L.L., L.N. Egorova, N.I. Rekhina, and M.N. Ereemeeva. 1956
Preservation (by acidification) of fish waste products and poor-quality fish
Rybn. Khoz. 32(9):78-83
- 252 Lagunov, L.L., L.N. Egorova, N.I. Rekhina, and M.N. Ereemeeva. 1958
Investigation of the process of acid preservation of fish and fish waste products
Tr. Vses. Nauch.-Issled. Inst. Morsk. Rybn. Khoz. Okeanogr. 35:115-130
- 253 Lambeth, J. 1970
Pyloric obstruction due to formic acid ingestion
Med. J. Malaya 24:187-189
- 254 Landes, D.R., and L.E. Dawson. 1971
Organic acid patterns in fresh and incubated whole egg products
Poult. Sci. 50(2):641-642
- * 255 Langler, J.E., and E.A. Day. 1966
Quantitative analysis of the major free fatty acids in Swiss cheese
J. Dairy Sci. 49(1):91-93
- 256 Langner, H.J. 1966
Aroma constituents of liquid smoke and smoking salts for the secondary smoking of foods
Fleischwirtschaft 46(9):961-966
- 257 Lanigan, G.W., and R.B. Jackson. 1965
Quantitative gas-liquid chromatography of volatile fatty acids. A method for the determination of C1 to C6 acids in biological material
J. Chromatogr. 17(2):238-244
- 258 Larsson, Ulla Britt, Torild Isaksson, and Olaf Samuelson. 1966
Anion exchange chromatography of organic acids in zinc acetate medium
Acta Chem. Scand. 20(7):1965-1969
- 259 Lee, Frank A., and R.S. Shallenberger. 1959
The flavor of canned peas in relation to their pyridolonecarboxylic acid content and to certain other factors
Food Res. 24(1):68-71
- * 260 Lehmann, G., and I. Lutz. 1970
Application of polyamide powder to analysis of organic preservatives
Z. Lebensmittelunters. Forsch. 144(5):318-321
- 261 Lemeshek, K. 1969
Chemical substances for foodstuffs conversation
Khodorovskaya 103 pp
- 262 Lenoel, C. Peaud. 1948
Determination in mixtures containing other volatile acids
Ann. Inst. Pasteur 74:345-346
- 263 Leopold, Heinrich, and Zdenek Valtr. 1959
Control of bacterial and yeast infections in citric acid fermentation
Czech. Pat. 91,768 issued Sep. 15, 1959
- 264 Jepper, H.A., M.T. Bartram, and Fred Hillig. 1956
Authentic packs of edible and inedible frozen eggs and their organoleptic, bacteriological, and chemical examination
J. Assoc. Offic. Agric. Chem. 39:185-194

- 265 Lin, Y.D., F.M. Clydesdale, and F.J. Francis. 1970
Organic acid profiles of thermally processed spinach puree
J. Food Sci. 35(5):641-644
- 266 Lindqvist, B., T. Storgards, and Maj Britt Goransson. 1953
Determination of the volatile acids in cheese by paper chromatography
Int. Dairy Congr. Proc. 13th 3:1250-1253
- 267 Lindsay, R.C., E.A. Day, and W.E. Sandine. 1965
Identification of volatile flavor components of butter culture
J. Dairy Sci. 48(12):1566-1574
- 268 Lipinska, Ewa, Marta Strzalkowska, Wilhelmina Goettlich, and W. Soltys. 1962
Application of nisin-resisting starters to the production of Edam cheeses
Int. Dairy Congr., Proc. 16th, Copenhagen, 1962 2:849-860
- 269 Luers, Heinrich, and Franz Weinfurter. No date
Über die wirksamkeitsbestimmung gewerblicher desinfektionsmittel
Wchschr. Brau. 43:25-29, 35-39, 45-49
- * 270 Lund, A. 1948
Excretion of methanol and formic acid in man after methanol consumption
Acta Pharmacol. Toxicol. 4:205-212
- * 271 Lund, A. 1948
Metabolism of methanol and formic acid in dogs
Acta Pharmacol. Toxicol. 4(2):108-121
- * 272 Lund, Alf. 1948
Metabolism of methanol and formic acid in rabbits
Acta Pharmacol. Toxicol. 4:99-107
- 273 Maeda, Yasuhiko. 1964
Organic acids in rhubarb
Nippon Shokuhin Kogyo Gakkaishi 11(10):444-446
- Maekawa, Kazuyuki, Masanobu Kodama, Mitsuo Kushi, Toshimitsu Mitamura, Nobuhisa Yada, and Hirotake Fukino. 1967
Gas chromatography of natural compounds
Mem. Ehime Univ., Sect. VI 12(3):131-147
- * 275 Malorny, G. 1959
The acute and chronic toxicity of formic acid and formates
Z. Ernährungswiss. 9(4):332-339
- 276 Malorny, G., and M. Rietbrock. 1962
Diet-dependent excretion of formaldehyde and formic acid in the urine of cats
Naturwissenschaften 49:520
- * 277 Malorny, Guenter. 1969
Metabolism studies with sodium formate and formic acid in man
Z. Ernährungswiss. 9(4):340-348
- 278 Malten, H. 1924
Generalized psoriasis following injection of formic acid
Munchen. Med. Wchschr. 70:336
- 279 Manabe, T. 1969
Studies on the constituents of and processing methods for Japanese chestnut. V. Organic acids in the edible portion of Japanese chestnut
J. Food Sci. Technol. 16(2):81-86
- 280 Mardashev, S.R., and L.A. Semina. 1948
Arginase in the skin
Biokhimiya 13:216-243
- 281 Marinetti, G.V., and E.R.M. Kay. 1963
The incorporation of C-14-formate into Ehrlich ascites tumor lipids
Biochim. Biophys. Acta 70:168-175
- 282 Markova, J. 1970
Qualitative and quantitative analysis of organic acids separated from rye flour, pre-ferment, dough and mix bread
Sh. Vysoke Skoly Chem.-Technol. Praze, E-Potravin 29:123-130
- 283 Matasa, C. 1965
New antifoaming agents used in the food industry
Ind. Aliment. 16(7):360-364
- 284 Mather, Alan, and Angel Assimos. 1965
Evaluation of gas-liquid chromatography in assays for blood volatiles
Clin. Chem. 11(11):1023-1035
- 285 Matsui, Harujiro, and Seichi Sato. 1963
Organic acids in poor sake
Nippon Jozo Kyokai Zasshi 58(7):635-638
- 286 Mattick, Leonard R., and Willard B. Robinson. 1960
Changes in the volatile acids during the baking of sherry wine by the Tressler baking process
Am. J. Enol. Viticult. 11:113-116
- 287 Mazur, M., S. Dzialek, J. Lembke, and H. Cichowicz. 1967
Effect of methanol, its metabolites, and some activators and inhibitors of carbohydrate metabolism on the isolated rabbit ileum
Agressologie 8(6):551-557
- 288 Mazur, Mieczyslaw, Stefan Dzialek, J. Lembke, and W. Dologicz. 1969
Effect of methanol and its metabolites on the contraction amplitude and coronary output of the isolated rabbit heart
Agressologie 10(4):317-324
- * 289 McInnes, A.G., R.P. Hansen, and A.S. Jessop. 1956
The volatile acids of mutton fat
Biochem. J. 63(4):702-704
- 290 McNeal, M.D., and C.J. Eldridge. 1922
Presence of formic acid in urine of infants and older children
Am. J. Dis. Child. 23:419-422
- 291 Mohlitz, A., and Th. Minas. 1965
Bitter orange peel oil. II. Terpenoids (oxygen-containing compounds)
Riechst. Aromen Koerperpflegen. 15(6):204-205, 207, 209, 211
- * 292 Meiselas, L.E. and A.A. Schlecker. 1957
The effect of nyloxin on the pain of arthritis
N.Y. State Med. J. 57(12):2067-2068
- 293 Merlis, M.N. 1939
Separation of formic acid from vinegar
Russ. Pat. 54,814 issued Apr. 30, 1939
- 294 Miler, M. 1931
Die analyse von fruchtessenzen
Russ.-Ger. Westnik Nauki Tech. 9:40-48
- * 295 Miller, P.G., P.L. Zimmerman, and E.B. Oberg. 1948
Comparison of vacuum and steam distillation for determining the volatile acidity of evaporated milk
J. Dairy Sci. 31(3):189-198
- 296 Minarik, E. 1971
Correction of the acidity of malts and wines
Bull. Office Int. Vin 44(479):57-63
- 297 Misler, Samuel, Leon Tumerman, and Henry R. Roberts. 1958
Distribution of carbon-14 in components of goat milk, with sodium formate-C-14 as precursor
Nature 181:346-347
- 298 Miyahara, Shojiro. 1961
Gas-liquid chromatographic separation and determination of volatile fatty acids in fish meat during spoilage
Nippon Suisan Gakkaishi 27:42-47

- 299 Mori, Takashi, Nobuo Muraoka, Kazuo Osodo, and Hanao Shitomi. 1966
Changes in organic acids of white peaches during postharvest ripening and cold storage
Nippon Shokuhin Kogyo Gakkaishi 13(9):375-380
- 300 Moriguchi, Shigehiro, and Hiroshi Ishikawa. 1964
Organic acids in soy sauce. VI. Changes in amounts of organic acids during the koji-making process
Hakko Kogaku Zasshi 42(1):35-40
- 301 Morin, Robert J. 1969
Incorporation of formate-C-14, acetate-1-C-14, choline-1,2-C-14, and ethanolamine-1,2-C-14 into rabbit ovarian phospholipids
Life Sci. 8(12):613-616
- 302 Moriyama, Shigetaka, Toshikatsu Ishiwara, and Yoshio Sakabe. 1964
Putrefaction of foods. IV. Study on putrefaction products of fish stored in a refrigerator, a hot box, in a thermostat, and at room temperature
Shokuhin Eiseigaku Zasshi 5(5):361-365
- * 303 Morr, C.V., W.J. Harper, and I.A. Gould. 1957
Some organic acids in raw and heated skim milk
J. Dairy Sci. 40(8):964-972
- 304 Morris, Mark M. 1935
Fluchtige sauren von wein
Ind. Eng. Chem. 27:1250-1252
- 305 Mosca, Americo. 1954
Preserving chocolate and its substitutes
Ital. Pat. 494,153 issued May 17, 1954
- 306 Motegi, S., and M. Matsubara. 1970
Causative bacteria of flat souring deterioration of fish sausage
J. Food Hyg. Soc. Jap. 11(5):369-372
- 307 Mukherjee, Kanai Lal, and Charles Heidelberger. 1962
Fluorinated pyrimidines. XV. Inhibition of the incorporation of formate-C-14 into deoxyribonucleic acid (DNA) thymine of Ehrlich ascites carcinoma cells by 5-fluoro-2'-deoxyuridine-5'-monophosphate and related compounds
Cancer Res. 22:815-822
- 308 Murphree, H.B. 1966
Effect of congeners in alcoholic beverages on the incidence of nystagmus
Quart. J. Stud. Alcohol. 27:201-213
- 309 Muthu, M., S.A. Jaleel, and V.S. Govindarajan. 1968
Storage of ripe areca fruits (Areca catechu): cold sterilization of fresh areca fruits with fumigants for prolonged storage
Trop. Sci. 10(3):160-166
- 310 Narasimhan, K.S. 1968
Fumigants as sterilants
Pestic., Symp. 1964:138-143
- 311 Nedelkovits, Janos, and Olga Szalay. 1962
Reaction mechanism of roasting of coffee surrogates. IV. Changes in the organic acid content of chicory during roasting
Budapesti Muszaki Egyet. Elelm-kem. Tansz. Közlem. 1962:27-33
- 312 Nohring, K., and I. Schroder. 1960
Preservation of animal waste substances. II. Preservation of blood from slaughterhouse animals with potassium metabisulfite or formic acid
Mh. Veterinarmed. 15:124-128
- 313 Nelson, E.K., and C.A. Greenleaf. 1929
Eine untersuchung von zuckerrohrsirupdestillations-ubertrieb
Ind. Eng. Chem. 21:857-859
- 314 Nelson, E.K., and H.H. Mottern. 1931
Some organic acids in honey
Ind. Eng. Chem. 23(3):335-336
- * 315 Neumark, H. 1967
On the areas of the stomach of sheep that are sensitive to formic acid and histamine
J. Agr. Sci. 69(2):297-303
- 316 Neumark, H., and A. Tadmor. 1968
The effect of histamine combined with formic or acetic acid on food intake and rumen motility, when infused into the omasum of a ram
J. Agric. Sci. 71(2):267-270
- 317 Neumark, H., et al. 1964
Amines, aldehydes, and keto-acids in silages and the effect on food intake by ruminants
J. Sci. Food Agr. 15:487-492
- 318 Nikiforova, V.N. 1958
Paper-chromatographic investigation of changes in sugars caused by heating
Tr. Vses. Nauch.-Issled. Inst. Konditer. Prom. 1958(13):20-34
- 319 Nixon, J.C., and S.H. Zbarsky. 1963
The effect of some transplantable tumors on the nucleic acid metabolism of the host tissues
Can. J. Biochem. Physiol. 41(11):2237-2247
- * 320 Noronha, J.M. and A. Sreenivasen. 1959
Formate metabolism in the vitamin B12 deficient rat
Biochem. J. 73:732-735
- 321 O'Rourke, P.J. 1950
Formic acid production amongst the formicidae
Ann. Entomol. Soc. America 43(3):437-443
- 322 Obata, Yataro, and Kagenori Matano. 1952
The flavor of katsuobushi. II. Identification of volatile substances by paper chromatography
J. Agric. Chem. Soc. Jap. 26:184-185
- 323 Obata, Yataro, and Kagenori Matano. 1953
The odor of cuttlefish-liver oil. I. Identification of volatile acids and bases
J. Oil Chem. Soc. (Japan) 2:112-114
- 324 Ockerman, Herbert W., Thomas N. Blumer, and H. Bradford Craig. 1964
Volatile chemical compounds in dry-cured hams
J. Food Sci. 29(2):123-129
- 325 Okada, H., S. Kashiwabara, and T. Sugiyama. 1939
Alkoholherstellung aus kartoffeln. IV. Über die maischebereitung mittels saurehydrolyse
Bull. Agric. Chem. Soc. (Japan) 14:10-11
- 326 Oldfield, J.F.T. 1957
Formation of volatile organic acids in factory juices and the estimation of these acids by gas-liquid chromatography
Compt. Rend. Assem. Comm. Int. Tech. Sucrerie, 10, London 1957:56-66
- 327 Olmsted, W.H., W.M. Whitaker, and C.W. Duden. 1929
Steam distillation of the lower volatile fatty acids from a saturated salt solution
J. Biol. Chem. 85(1):109-114
- 328 Olsson, Nils. 1942
The preservation of fish products and their suitability as feed for hens and chicks
Lantbrukshogskol., Husdjursforsoksanst., Medd. 7:1-55
- 329 Olsson, Nils, and Olofsson, N. Edvin. 1955
The preservation of fish products for feeding
Kgl. Lantbrukshogskol. Och Statens Lantbruksforsok, Statens Husdjursforsok Medd. 59:38 pp
- 330 Opienska-Blauth, Janina, Olga Saklowska-Szymonowa, and Marek Kanski. 1951
Chromatografia bibulowa niektorych kwasow organicznych. (The partition paper chromatography of some organic acids)
Ann. Univ. Mariae Curie-Sklodowska, Sect. D-Med. 5(9):221-275

- * 331 Oro, J. and D.A. Rappoport. 1959
Formate metabolism in animal tissues. II. The mechanism of formate oxidation
J. Biol. Chem. 234(7):1661-1665
- 332 Oro, John F., and D.A. Rappoport. 1957
Formate metabolism by animal tissues. I. Metabolism of formate-C-14 by isolated rabbit and rat jejunum
J. Biol. Chem. 224(1):489-498
- 333 Oshmyan, G.L., and A.V. Ignatova. 1961
Chemistry of the action of activated charcoal upon alcohols
Tr., Tsent. Nauchn.-Issled. Inst. Spirt. Likero-Vodochn. Prom. 11:242-250
- 334 Osman, H.G., and Afaf El-Mariah. 1960
Studies on the inhibitory effect of combined chemical preservatives on *Saccharomyces cerevisiae*
J. Am. Pharm. Assoc., Pract. Pharm. Ed. 49:231-233
- 335 Ostroumova, M.N. 1966
Inclusion of formic acid and single-carbon fragments of serine and histidine in purines of nucleotides and nucleic acids from mice hematoma
Nukleinovye Kisloty, Tr. Konf., 2nd 1965:315-318
- 336 Pace, Donald M., Byron Th. Aftonomos, Alice Elliott, and Steven Sommer. 1967
Effects of the sodium salts of certain monocarboxylic acids on established cell lines
Can. J. Biochem. 45(1):81-88
- 337 Palloy, E., E. Sundt, and M. Winter. 1963
Aromas. VII. Analysis of the aroma of raspberries. 1. Lower acids and esters
Helv. Chim. Acta 46(6):2297-2299
- 338 Patton, S., and R.J. Flipse. 1957
Carbon-14 activity of some heat-degradation products of milk containing lactose-1-C-14
Science 125:1087-1088
- 339 Paulose, M.M., and K.T. Achaya. 1964
Stability of castor seeds and oil. II. Nature of lipase action and of hull oil
Indian Oilseeds J. 8(4):324-327
- 340 Poud Lenoel, C. 1948
Dosage de l'acide formique en melange avec d'autres acides volatils
Ann. Inst. Pasteur (Paris) 74(4):345-346
- 341 Peck, Harry D., Jr., and Howard Gest. 1957
Formic dehydrogenase and the hydrogenase enzyme complex in *coli-aerogenes* bacteria
J. Bact. 73(6):706-721
- 342 Penczek, Stanislaw. 1960
Investigation of progress in synthesis of epoxy-type plasticizers
Przem. Chem. 39:232-236
- 343 Penfold, A.R. No date
Die keimtotenden eigenschaften einiger australischer atherischer ole und ihrer reinen bestandteile
Perfum. Essent. Oil Record 17:251-253
- 344 Perila, Olavi. 1956
Saturated fatty acids of stem cells or firch (*Betula verrucosa*) and aspen (*Populus tremula*)
Ann. Acad. Sci. Fenn. Ser. A. II. Chem. 76:7-49
- 345 Perila, Olavi. 1956
Separation of saturated straight-chain fatty acids III. Quantitative paper chromatography
Acta Chem. Scand. 10:143-144
- 46 Perretta, Marco, and Maria Pieber-Perretta. 1962
Species difference in the pattern of formate incorporation into marrow nucleic acids in vitro
Biochim. Biophys. Acta 61:828-829
- 347 Perretta, Marco, Wilhelm Rudolph, Gloria Aguirre, and George Hodgson. 1964
Uptake of formate by bone marrow and spleen nucleic acids in normal and erythrocyte-transfused mice
Biochim. Biophys. Acta 91(2):335-337
- 348 Perretta, Marco, Wilhelm Rudolph, Gloria Aguirre, and George Hodgson. 1964
Metabolic pattern of formate-C-14 incorporation in vivo and in vitro into the nucleic acids of normal mouse spleen and bone marrow
Biochem. Biophys. Acta 87(1):157-159
- 349 Peyron, Louis. 1962
Study of essences from liquors obtained by the steam distillation of Morocco rose geranium
Compt. Rend. 255:2981-2982
- 350 Pickett, M.J., H.L. Ley, and N.S. Zygmuntowicz. 1944
Manometric determination of formic acid
J. Biol. Chem. 156(1):303-315
- 351 Pittarelli, M. 1935
New applications for old reaction for determination in organic fluids
Minerva Med. 1:428-429
- 352 Platon, R., A. Orborn, and K.E. Thome. 1944
Acidulation with acetic or formic acid of skim milk used for feeding animals
Svenska Mejeritidn. 36:255-262
- 353 Pohloudek-Pabini, R., and K. Luthardt. 1958
Paperchromatographic detection of esters of lower fatty acids in essential oils
Naturwissenschaften 45:314
- 354 Poiger, H., G. Falkner, and H. Michl. 1969
Enzymic determination of formate with formic acid dehydrogenase from peas
Mikrochim. Acta 1969(6):1165-1169
- 355 Pontovich, V.E., and L.M. Pinchuk. 1958
Content of volatile acids in the fruit of oily poppy
Dokl. Akad. Nauk S.S.S.R. 120:585-587
- 356 Poore, H.D. 1932
Zusammensetzung von californischen citronen-und pomeranzenolen
Perfum. Essent. Oil Record 23:166-172
- 357 Popovskii, V.G., O.A. Timofeeva, and I.M. Soboleva. 1968
Alteration in aromatic substances and the color of fruits by sublimation drying during preservation (Russian)
Tr. Mold. Nauch.-Issled. Inst. Pishch. Prom. 8:70-81
- 358 Power, Frederick B., and Victor K. Chesnut. 1921
Die wohlriechenden bestandteile der pfirsiche
J. Am. Chem. Soc. 43:1725-1739
- 359 Pribela, A., and P. Rohanek. 1965
Changes taking place in aromatic substances during fruit sirup manufacture by various methods
Prun. Potravin 16(4):199-201
- 360 Prodanski, P. 1964
Some volatile aromatic components of cheese
Prun. Potravin. 15:273-274
- 361 Puskarevs, I. 1965
Chromatographic method for quantitative determination of volatile fatty acids in biological samples. Volatile fatty acids in perienteric fluid of *Ascaris suum*
Latv. PSR Zinat. Akad. Vestis 1965(3):93-99
- 362 Quijcho, Florante A., and Felix Friedberg. 1962
Action of concentrated acids on adenosine triphosphate (ATP)-creatine phosphotransferase
Can. J. Chem. 40:1018-1022

- * 363 Rabinowitz, Jesse C., and Herbert Tabor. 1958
The urinary excretion of formic acid and
formamino-glutamic acid in folic acid deficiency
J. Biol. Chem. 233(1):252-255
- 364 Radtke, R., W. Mohr, and R. Springer. 1966
Coffee aromas and their changes during storage of
roasted coffee. III. Determination of low-
volatile aroma compounds and volatile acids
Z. Lebensm.-Unters.-Forsch. 129(6):349-359
- 365 Raible, Karl. 1957
Effect of age of yeast cultures on the
effectiveness of chemical preservatives
Deut. Lebensm.-Rundschau. 53:198-200
- 366 Raible, Karl. 1957
Effect of the initial concentration of yeast
cells on the effectiveness of
chemical preservatives in malt-wort
Deut. Lebensm. Rundschau. 53:155-156
- 367 Raible, Karl. 1957
Effect of sub-threshold concentrations of
chemical preservatives on the growth
and fermentation of yeast in maltwort
Dtsch. Lebensm.-Rundsch. 53:60-64
- 368 Raible, Karl, and Gisela Busch. 1957
Studies on chemical conserving materials. I. The
influence of the common preservatives on the
multiplication of yeasts
Zentr. Bakt., Parasitenk., Abt. II 110:172-177
- 369 Ralls, Jack W. 1960
Flash exchange gas chromatography for the
analysis of potential flavor components of peas
J. Agric. Food Chem. 8:141-143
- * 370 Rappoport, D.A., B.W. Sewell and F.B. Moreland.
1959
Synthesis of proteins, lipids and nucleic acids
from formate in young rats
Am. J. Physiol. 197(1):1045-1047
- 371 Rappoport, Donald A., James A. Green, and Joseph
H. Gast. 1956
Formate oxidation by erythrocytes
Arch. Biochem. Biophys. 63(2):343-351
- 372 Rastelli, G., and C. Gibelli. 1930
Sul principio attivo dello Xanthium spinosum
(Active principle of X. spinosum)
Boll. Soc. Ital. Biol. Sper. 5(4):549-554
- 373 Ravenx, Roger, and Josy Bove. 1957
Separation and determination of carboxylic acids
from C1 to C6 by partition chromatography on a
silica column. I. General Study
Bull. Soc. Chim. Fr. 1957:363-376
- 374 Roche, O.R. 1967
Analysis of food preservatives presently allowed
in Germany
Z. Lebensm.-Unters. Forsch. 133(6):375-386
- 375 Recke, Wilhelm. 1934
Untersuchungen über die kochentechnische
verwendbarkeit von kochsalzersatzpräparaten
Arch. Verdauungskr. 55:274-281
- 376 Rehm, H.J. 1963
Influence of chemical compounds on the
antimicrobial preservatives. I. Influence of
various types of compounds on the preservative
action against Aspergillus niger
Z. Lebensm.-Unters.-Forsch. 118:508-522
- 377 Rehm, H.J., F. Senner, and E.M. Lukas. 1964
Action of preservative combinations. IX. Action
of binary and ternary combinations
of preservatives with nisin and tylosin
Z. Lebensm.-Unters.-Forsch. 125(4):258-271
- 378 Reinhold, J., H. Lucas, and R. v. Veh. 1943
The "hard cooking" of naturally soured vegetables
Gartenbauwiss. 18:1-26
- 379 Rety, J. 1930
Dermatitis provoked by action of formates and a
surface-active agent
Rev. Lyon Med. 15:761-762
- 380 Ricceri, G., and A.M. Giuffrida. 1961
Incorporation of formate-C-14 into bases of the
nucleic acids of organs of rats fed with zein as
the sole amino acid source
Riv. Sedute Accad. Giochia Sci. Nat. Catania
6:185-199
- 381 Rice, A.C., and C.S. Pederson. 1954
Factors influencing growth of Bacillus coagulans
in canned tomato juice. II. Acidic constituents
of tomato juice and specific organic acids
Food Res. 19(1):124-133
- 382 Rice, A.C., and C.S. Pederson. 1954
Chromatographic analysis of organic acid in
canned tomato juice, including
the identification of pyrrolidone-carboxylic acid
Food Res. 19(1):106-114
- 383 Rice, Andrew C., and Carl S. Pederson. 1954
Factors influencing growth of Bacillus coagulans
in canned tomato juice. I. Size of inoculum and
oxygen concentration
Food Res. 19:115-133
- 384 Richards, E.L., and M.R. Chandrasekhara. 1960
Chemical changes in dried skim milk during storage
J. Dairy Res. 27:59-66
- 385 Riedler, G., et al. 1968
Effects of acute intravascular hemolysis on
coagulation and fibrinolysis. 3. Hemoglobinemia
caused by ingesting
Helv. Med. Acta 34:223-230
- 386 Riethbrock, Norbert, and Wolf Dietrich Hinrichs.
1964
A simple method for the quantitative
determination of formic acid in human urine and
blood
Klin. Wochschr. 42(19):981-985
- 387 Ritchie, E.B. and W.C. King. 1953
Observations on the reaction of larvae of
ancylostoma canium and a. braziliense to various
drugs and chemical compounds based on in vitro
tests
J. Invest. Dermatol. 20(5):337-341
- 388 Ritter, W., and H. Haenni. 1960
Gas chromatography and its application in the
dairy industry. II. Detection and determination
of volatile fatty acids in dairy products and
cultures
Milchwissenschaft 15(6):296-302
- 389 Roe, F.J.C. 1955
Tumour-initiating action of urethane and its
inhibition by purine precursors
Nature 175(4458):636-637
- 390 Roe, Oluf. 1946
Methanol poisoning. Its clinical course,
pathogenesis, and treatment
Acta Med. Scand. 126(Suppl. 182):253 pp
- 391 Rogers, Terence A., and Max Kleiber. 1956
The biosynthesis of milk-fat in the intact cow
Biochim. Biophys. Acta 22(2):284-292
- 392 Rosculet, George. 1969
Metabolic products of yeast fermentation: organic
acids, isolation, separation,
and characterization
Am. Soc. Brew. Chem., Proc. 1969:127-138
- 393 Rostkowska, Julia. 1964
Effects of chemical agents on the motor responses
in Balantidium coli
Acta Protozool. 2(1-9):81-89

- 394 Rother, H. 1961
Estimation of preservatives in fruit juices and concentrates
Ind. Obst. Gemuseverwert. 46:638-642
- 395 Rubashkina, S.Sh., L.L. Kukhar'kova, and P.V. Perova. 1959
The effectiveness of various preservatives in the processing of blood for food purposes
Tr. Vses. Nauch.-Issled. Inst. Myasnoi Prom. 1959(9):75-79
- 396 Rubinstein, Riva, Ann V. Howard, and O.M. Wrong. 1969
In vivo dialysis of feces as a method of stool analysis. IV. Organic anion component
Clin. Sci. 37(2):549-564
- 397 Ryss, Simon. 1932
Über die Wirkung des Ameisensauren Natriums und einiger anderer Natronsalze auf die Magendrusensekretion
Z. Ges. Exp. Med. 82:382-386
- 398 Sacharow, I.P. 1937
Die Acidität des Mehls
Chem. J. Ser. B. J. Angew. Chem. 10:1095-1101
- 399 Sakami, Warwick, and Arnold D. Welch. 1950
Synthesis of labile methyl groups by the rat
J. Biol. Chem. 187:379-384
- 400 Sammartino, U. 1933
Comparative symptomatology of minimal lethal doses of methyl alcohol, formaldehyde and formic acid
Arch. Pharmacol. Sper. 56:351-363
- 401 Sammartino, U. 1933
Comparative action of methyl alcohol, formaldehyde and formic acid on posterior circulation of frog
Arch. Pharmacol. Sper. 56:320-326
- 402 Sammartino, U. 1933
Comparative action of methyl alcohol, formaldehyde and formic acid on isolated frog's heart
Arch. Pharmacol. Sper. 56:311-319
- * 403 Sammartino, U. 1933
Research on the toxicity of methyl alcohol. More rational estimate and interpretation of the mode of action of methyl alcohol.
Arch. Pharmacol. Sper. 56:364-371
- 404 Sandino, W.E., and P.R. Elliker. 1969
Flavour in fermented dairy products
Abstracts of Papers. Am. Chem. Soc. 158:MICR 29
- 405 Sarris, J., J.N. Morfaux, P. Dupuy, and D. Hertzog. 1970
Automatic determination of volatile acidity in wine
Ind. Aliment. Agric. 87(2):115-121
- * 406 Sarudi, I., Jr. 1969
Modified steam distillation procedure for the estimation of formic acid in fruit syrups
Naturbrunnen 19(3):106, 108
- 407 Sarudi, I., Jr. 1969
Modified steam distillation method for the determination of formic acid in fruit syrups
Konzerv-os Paprikaipar 1969(2):32-34
- 408 Schneider, P. 1928
Poisoning; case
Beitr. Gerichtl. Med. 8:212-216
- 409 Schreier, M. 1966
Experiments with organic acids for the use as disinfectants in the meat-processing industry
Mh. Veterinarmed. 21:738-748
- 410 Schroder, I., and W. Laube. 1960
Chemical preservation of fish wastes and food fishes
Deut. Landwirtschaft. 11:245-247
- 411 Schweigart, H.A. 1969/1970
The Biotest. Development of a biological wholesomeness test
Vitalst.-Zivilisationskr. 14(1):19-21, (2):61-64, (3) 101-103, (4):141-144, (5):191-193
- 412 Schweisheimer, W. 1955
Extraction of Californian orange-oil: the odor principle in orange skins
J. Sci. Food Agric. 7:1-236
- 413 Scott, T.W., I.G. White, and E.F. Annison. 1962
Oxidation of short-chain fatty acids by ram, bull, dog, and fowl spermatozoa
Biochem. J. 83:392-398
- * 414 Sedlacek, B.A.J., and St. Prochazka. 1954
Content of formic acid in fruit and fruit products
Prun. Potravin 5:491-493
- 415 Semichon, L., and M. Planzy. 1932
Untersuchung über die organischen Säuren der weine, traubennoste und natürlichen oder vergorenen fruchtsäfte. II. Die flüchtigen Fettsäuren
Ann. Sci. Agron. Fr. 2:504-528
- 416 Senczek, D. 1972
Storage of delicatessen salads
Feinkostwirtschaft 9(1):13-16
- 417 Shaburov, M.A., V.F. Demarina, and K.L. Tsekanskaya. 1970
Chromatographic determination of low-molecular-weight monocarboxylic acids and their butyl esters
Neftepererab. Neftekhim. 1970(4):46-48
- 418 Shah, N.C., and B.C. Gulati. 1965
Water-soluble constituents of palmarosa oil
Cymbopogon martinii var. motia
Perfum. Essent. Oil Record 56(7):435-437
- 419 Shaposhnikov, V.N., and Zh. K. Loriya. 1967
Pathways of utilization of formic acid by Bacterium formicum
Dokl. Akad. Nauk. S.S.S.R. 175(3):721-722
- * 420 Shelley, Ruth N., Harold Salwin, and William Horwitz. 1963
Quantitative determination of formic, acetic, propionic and butyric-acids by gas chromatography (in food technology)
J. Assoc. Offic. Agric. Chem. 46(3):486-493
- 421 Shifrina, Kh. B., T.P. Dvornikova, N.N. Zaginailo, Ya. N. Kazanovich, and K.D. Shchupak. 1963
Biochemistry of tomatoes
Biokhim. Kul't. Rast., Akad. Nauk Moldavsk. SSR, Inst. Fiziol. Biokhim. Rast., Moldavsk. Nauch.-Issled. Inst. Oroschaemogo Zemled. Ovoshchevodstva 3:3-69
- 422 Simmons, Perez, and C.K. Fisher. 1945
Ethyl formate and isopropyl formate as fumigants for packages of dried fruits
J. Econ. Entomol. 38:715-716
- 423 Simonart, Paul, and Jacques Mayaudon. 1956
Chromatographic study of cheese. II. Aliphatic acids (French)
Neth. Milk Dairy J. 10:156-161
- 424 Smellie, R.M.S., W.M. McIndoe, and J.N. Davidson. 1953
Incorporation of nitrogen-15, sulfur-35, and carbon-14 into nucleic acids and proteins of rat liver
Biochim. Biophys. Acta 11:559-565
- * 425 Smillie, L.B. and H. Neurath. 1959
Reversible inactivation of trypsin by anhydrous formic acid
J. Biol. Chem. 234(2):355-359

- 426 Smirnova, G.M., and L.A. Egorova. 1963
Effect of enzymic preparations on the formation
of volatile acids in rye-leaven and dough
Tr. Tsent. Nauchn.-Issled. Inst. Khlebopekarn.
Prom. 1963(9):109-114
- 427 Smith, D.E., and J.R. Coffman. 1960
Analysis of food flavors by gas-liquid
chromatography. Separation and identification of
the neutral components from bread preferment
liquid
Anal. Chem. 32:1733-1737
- 428 Smouse, Thomas H., and Stephen S. Chang. 1967
A systematic characterization of the reversion
flavor of soybean oil
J. Am. Oil Chem. Soc. 44(8):509-514
- * 429 Smyth, H.F., C.P. Carpenter, C.S. Weil and U.C.
Pozzani. 1954
Range-finding toxicity data. List V
Arch. Ind. Hyg. Occup. Med. 10:61-68
- * 430 Sollmann, T. 1921
Studies of chronic intoxications on albino rats;
acetic and formic acids
J. Pharm. Exp. Ther. 16:463
- 431 Sone, Kiyohide. 1962
Inhibition of amylase activity by flavines. II.
Inactivation of alpha-amylase by fluorescent
substances
Nippon Nogeikagaku Kaishi 36(1):7-12
- 432 Spanyol, P., and I. Szeredy. 1962
Smoking of foods. IV. Available acids in smoke
and smoked products
Z. Lebensm. Unters. Forsch. 118:293-299
- * 433 Spector, W.S., (Ed.). 1956
Handbook of Toxicology; Vol. I, Acute Toxicities
W. B. Saunders Company, Philadelphia
- 434 Sperling, P., E.S. Maxwell and W.P. von Oettingen.
1953
Comparative excretion and distribution of C14-
labeled carbonate and formate in large albino rats
Am. J. Physiol. 174:33-38
- * 435 Sporn, A., Victoria Marin, and C. Shoebesch. 1962
Toxicity of formic acid
Ighena (Bucharest) 11:507-515
- 436 Standard Brands Inc., New York. 1936
Verhinderung von schimmelbildung auf brot
E. Pat. 455,846 issued Nov. 26, 1936
- 437 Standard Oil Co. of California, San Francisco.
1935
Denaturierte athylester
U.S. Pat. 1,995,539 issued Mar. 26, 1935
- * 438 Stecher, P.G., (Ed.). 1968
The Merck Index-An Encyclopedia of Chemicals and
Drugs, 8th ed.
- 439 Stein, Abraham M., and John W. Mehl. 1958
The effect of implanted tumors on the oxidation
of formate and of the methyl group of methionine
in mouse liver
Cancer Res. 18:1196-1202
- 440 Stein, Abraham M., William T. Murakami, and
Donald W. Visser. 1959
Purine synthesis in mouse tissues
Cancer Res. 19:84-89
- 441 Steiner, P. 1928
Poisoning; case
Mitt. Gesellsch. Inn. Med. Kinderh. 27:39
- 442 Steinhauer, John Edward, and Lawrence E. Dawson.
1969
Quantitative determination of formic, acetic,
propionic, and butyric acids in frozen whole
eggs by gas-liquid chromatography
J. Food Sci. 34(4):359-364
- 443 Stepp, W. No date
Über den befund von ameisensäure im menschlichen
blute
Z. Physiol. Chem. 109:99-107
- 444 Stepp, Wilhelm, and Hermann Zumbusch. 1920
Studien über den intermediären
kohlenhydratstoffwechsel beim menschen. II. Über
das quantitative verhalten der ameisensäure im
normalen und pathologischen blute
Dtsch. Arch. Klin. Med. 134:112-118
- * 445 Stern, H. 1906
Sixteen years experience with formic acid as a
therapeutic agent
J. Am. Med. Assoc. 46:1258-1262
- 446 Stinson, Edgar E., Mary H. Subers, Jeanne Petty,
and Jonathan W. White, Jr. 1960
The composition of honey. V. Separation and
identification of the organic acids
Arch. Biochem. Biophys. 89:6-12
- * 447 Stokstad, E.L., R.E. Webb and E. Shah. 1966
Effect of vitamin B12 and folic acid on the
metabolism of formiminoglutamate, formate and
propionate in the rat
J. Nutr. 88:225-232
- 448 Storgards, Torsten, and Borge Lindqvist. 1952
Different methods for demonstrating faulty
fermentation in cheese
Svenska Mejeritidn. 44:103-107
- 449 Stumm-Tegethoff, B.F.A. 1964
Formaldehyde impurities as a mutagenic principle
in drosophila melanogaster
Naturwissenschaften 51:646-647
- * 450 Stumm-Tegethoff, B.F.A. 1969
Formaldehyde-induced mutations in drosophila
melanogaster in dependence of the presence of
acids
Theor. Appl. Genet. 39:330-334
- 451 Suzue, Mohei. 1927
Untersuchungen über die physikalisch-chemischen
eigenschaften der zellmembran
J. Biophys. 2:31-80
- 452 Suzuki, Taneko. 1953
Determination of volatile acids for judging the
freshness of fish
Bull. Jap. Soc. Sci. Fish. 19(2):102-105
- 453 Szonntag, E. 1959
Determination of the total and some more
important volatile acids in cheese
Int. Dairy Congr., Proc., 15th Congr., London
3:1467-1473
- 454 Tanaka, Kenji, Kazuyo Furukawa, and Hiroshi
Matsumoto. 1967
The effects of organic and inorganic acids on the
physical properties of dough
Hakko Kogaku Zasshi 45(6):566-569
- 455 Tanikawa, Eiichi, and Terushige Motohiro. 1960
Refrigeration of salmon as the raw material for
canning
Reito 35:15-23
- 456 Tarantola, C. 1971
Composition of grappa in relation to the
development of the volatile constituents of the
pressed grape lees during storage and to the
distillation technique
Vini Ital. 13(73):331-335
- 457 Tasman, Abel, and Arnold William Pot. 1935
Die bildung von wasserstoffgas aus traubenzucker
und ameisensäure durch sogenannte "ruhende
Colibacillen"
Biochem. J. 29:1749-1756
- 458 Taylor, Nelson W. 1917
Das eindringen von saure in lebende gewebe
J. Gen. Physiol. 11:207-219

- 459 Teramoto, S., H. Taguchi, and K. Ueda. 1962
Organic acid constituents of Japanese fermented
food products
Proc. Int. Congr. Food Sci. Technol., 1st,
London, 1962 2:549-555
- 460 Terashima, Hiroko. 1963
Changes of organic acids in salt-bran pickling
Eiyo To Shokuryo 16(4):249-253
- 461 Theis, Edwin R. 1930
Eine kritische studie zur biochemie des weichens
und ascherns. V. Der einfluss
verschiedener sauren auf die diffusion
koagulierbarer proteine aus tierischer haut
J. Am. Leather Chem. Assoc. 25:48-56
- 462 Thompson, Adrienne R. 1951
Volatile products of apples. I. Identification of
acids and alcohols
Austral. J. Sci. Res. B4:283-290
- 463 Toennissen, E., and E. Brinkmann. 1938
The catabolism of lower fatty acids, especially
acetic and formic acids, in mammals and the
formation of sugar from fat
Z. Physiol. Chem. 252:169-195
- * 464 Tomic-Nedeljkovic, M. 1970
Contents of methanol and formic acid in some
Yugoslav wines
Arch. Farm. 19(4):225-228
- 465 Tomizawa, Jun-ichi. 1954
On the mechanism of aconitase action. III.
Kinetic analysis using DL-isocitric acid-2-C14
J. Biochem. (Tokyo) 41(5):567-575
- 466 Tranger, Margareta. 1963
A method for quantitative determination of lower
fatty acids in serum by paper chromatography
Vet. Archiv. 33:248-252
- 7 Tucek, Arnost, and Vlastimil Sanda. 1951
Analytical determination of chemical preserving
agents in fruit and vegetable products
Chemie (Prague) 7:203-206
- 468 Turowska, Maria. 1960
Biacridine derivatives as indicators in titration
of weak acids
Chem. Anal. (Warsaw) 5:815-821
- * 469 U.S. Department of Commerce. 1972
U.S. Foreign Trade Imports. Commodity by Country
U.S. Department of Commerce, Washington, D.C.
PT 135
- 470 Ueda, Ryuzo, Shiro Nagai, and Shigehiro Moriguchi.
1959
Organic acids in soysauce. I. Determination of
the organic acids in soysauce
Hakko Kagaku Zasshi 37:94-99
- 471 Ueno, Takahiro. 1961
Semi-chemical soy sauce. IX. Inhibitory effects
of an acid hydrolyzate of defatted soybeans on
yeast fermentation
Nippon Nogei Kagaku Kaishi 35:1184-1189
- 472 Ueno, Takahiro, and Taira Kuramochi. 1961
Semi-chemical soy sauce. VII. Volatile components
produced by HCl hydrolysis of defatted soybean
Nippon Nogei Kagaku Kaishi 35:454-458
- 473 Umeh, Eric O. 1970
Separation and determination of low-molecular
weight straight-chain (C1-C9) carboxylic acids
by gas chromatography of their anilide derivatives
J. Chromatogr. 51(2):147-154
- 474 Union Carbide Corp. 1968
Toxicology studies: formic acid
Industrial Medicine and Toxicology Dept., Union
Carbide Corp. (data sheet)
- * 475 Usseglio-Tomasset, Luciano. 1967
Formic acid in wines
Accad. Ital. Vite Vino, Siena, Atti 19:229-242
- 476 Usvyatsov, A.A., and V.T. Solomatin. 1970
Amperometric titration of formaldehyde and formic
acid
Zavod. Lab. 36(2):154-155
- 477 V. Schelhorn, M. 1948
Activity and application of chemical preserving
agents in food industry
Dtsch. Lebensm.-Rundsch. 44:144
- 478 V. Schelhorn, M. 1951
Preservatives. V. Comparative effectiveness of
preservatives
Dtsch. Lebensm. Rundsch. 47:16-18
- 479 Van Dame, Halver C. 1954
Decomposition in tomato products. Determination
of acetic, formic, succinic, and lactic acids
J. Assoc. Offic. Agric. Chem. 37:572-574
- 480 Vashkov, V.I., and A.K. Astaf'eva. 1954
The virucidal and bactericidal action of some
organic acids
Tr. Tsentral. Nauch. Issled. Dezinfekts. Inst.
1954(8):27-30
- 481 Venkataraman, S., D.K. Biswas, and B. Connor
Johnson. 1967
Effect of propionate on the induction of vitamin
B12 deficiency in chicks and rats
J. Nutr. 93(2):131-134
- * 482 Vercoe, J.E., and K.L. Blaxter. 1965
The metabolism of formic acid in sheep
Brit. J. Nutr. 19(4):523-530
- 483 Veringa, H.A., T.E. Galesloot, and H. Davelaar.
1968
Symbiosis in yoghurt. II. Isolation and
identification of a growth factor
for Lactobacillus bulgaricus produced by
Streptococcus thermophilus
Neth. Milk Dairy J. 22(3):114-120
- 484 Villegas, Trinidad S. 1936
Quantitative relations of various organic
substances in the fruit of Sandoricum koetjabe
Univ. Philipp. Nat. Applied Sci. Bul. 5:449-450
- * 485 Vincent, L.E. and D.L. Lindgren. 1972
Hydrogen phosphide and ethyl formate: fumigation
of insects infesting dates and other dried fruits
J. Econ. Entomol. 65(6):1667-1669
- 486 Voit, K. 1928
Ueber das verhalten der ameisensaure im
diabetischen organismus
Z. Klin. Med. 109:227-268
- 487 Voit, K., and H. Friedrich. 1935
Appearance in course of apple diet
Klin. Wchnschr. 14:1742-1793
- 488 Voit, Kurt. 1928
Ueber das verhalten der ameisensaure im
diabetischen organismus. (The status of
formic acid in diabetics)
Z. Klin. Med. 106(3/4):227-268
- 489 Von Euler, Hans, Hans Hasselquist, and Alf Larsen.
1962
Cleavage of deoxyribonucleic acid (DNA) with
formic acid
Ark. Kem. 18:533-540
- 490 Von Fellenberg, Th. 1936
The determination of formic acid in fruit juice
and fruit sirup
Mitt. Lebensm. Hyg. 27:182-200
- * 491 Von Ottingen, W.P. 1959
The aliphatic acids and their esters--toxicity
and potential dangers. The saturated monobasic
aliphatic acids and their esters
Arch. Ind. Health 20(6):517-531

- 492 Von Schelhorn, M. 1953
Bacteriostatic and bactericidal action of preservatives
Arch. Mikrobiol. 19:30-44
- 493 Von Schelhorn, Mathilde. 1953
Efficacy and specificity of chemical food preservatives
Food Technol. 7:97-101
- 494 Von Szelenyi, Geza. 1932
Titrimetrische bestimmungsmethoden der ameisensaure in fruchtsaften und fruchtsirupen
Z. Unters. Lebensm. 63:534-541
- 495 Vrchlabsky, J. 1969
The influence of sodium metabisulphite in combination with sulphuric, hydrochloric or formic acid upon certain bacteria and viruses
Proc. Eur. Meet. Meat Res. Work. 15:193-198
- * 496 Waldo, D.R., J.F. Keys and C.H. Gordon. 1973
Preservation efficiency and dairy heifer response from unwilted formic acid, and wilted untreated silage
J. Dairy Sci. 56(1):129-136
- 497 Wallenstein, H.D. 1959
Organic acids in beets and juices. I. Volatile acids
Z. Zuckerind. 9:611-614
- 498 Wallenstein, H.D., and Heinz Egler. 1960
Organic acids in beets and juices. III. Application of the Craiy extractor to the determination of volatile acids
Z. Zuckerind. 10(2):66-71
- 499 Wallenstein, H.D., and K. Bohn. 1963
Organic acids in beets and juices. IV. Changes in anion content during sugar extraction process
Z. Zuckerind. 13(3):126-131
- 500 Walter, Erich. 1931
Die haltbarkeit der mit ameisensaure konservierten fruchtsäfte
Destillateur Likorfabrikant 44:548-550
- * 501 Walter, Erich. 1961
The natural formic acid content of fruit juices
Alkohol Ind. 1961(15):391-392
- 502 Wang, Kunq-Tsung. 1960
Polyamide-impregnated paper chromatography
J. Chin. Chem. Soc. 7:64-68
- * 503 Watson, J.R., and P. Crescuolo. 1970
Gas chromatographic separation and detection of C1 to C1 monocarboxylic acids as the p-substituted benzyl esters
J. Chromatogr. 52(1):61-67
- * 504 Watts, D.T. 1951
The effect of methanol, formaldehyde, and formic acid on tissue respiration
Arch. Biochem. Biophys. 34:391-395
- 505 Webb, A. Dinsmoor, Richard E. Kepner, and W.G. Giletto. 1964
Comparison of the aromas of flor sherry, baked sherry, and submerged-culture sherry
Am. J. Enol. Viticult. 15(1):1-10
- * 506 Wheeler, Glynn P., Jo Ann Alexander, and Harold P. Morris. 1964
Synthesis of purines and nucleic acids and catabolism of purines by rat liver and hepatomas
Adv. Enzyme Regul. 2:347-369
- 507 Wierzchowski, J., and H. Witkowski. 1966
Paper-chromatographic determination of short-chain volatile fatty acids of fish
Mitt. Geb. Lebensmittelunters. Hyg. 57(2):101-127
- 508 Wilhelm, H., F. Palluy, and M. Winter. 1966
Flavors. XII. The flavor of fresh strawberries. Identification of the volatile acids and other compounds (French)
Helv. Chim. Acta 49(1):65-67
- 509 Wilmanns, W. 1961
The tetrahydrofolic acid activation of one-carbon compounds in normal and pathologic white blood cells
Klin. Wochschr. 39:884-888
- 510 Windisch, Siegfried. 1958
Formation of volatile acids by pellicle-forming yeasts
Branntweinwirtschaft 80:21-22
- 511 Wohack, Franz. 1921
Mikroanalytische verfahren in der nahrungsmitteluntersuchung
Z. Unters. Nahrgrs. Genussm. 42:290-298
- 512 Wojtowicz, Maria. 1961
Spectrophotometric analysis of the fruit juices and sirups. I. The effect of certain physical and chemical agents
Zesz. Nauk. Szk. Gl. Gospod. Wiejsk. Warsz., Roln. 4:149-195
- 513 Wood, N.A. 1969
Formic acid as a prophylactic for candidasis in partridges
Vet. Rec. 85(4):78-80
- 514 Woodman, J.S., Andre Giddey, and Robert H. Egli. 1968
Carboxylic acids of brewed coffee
Int. Colloq. Chem. Coffee, 3rd 1967:137-143
- * 515 World Health Organization of the United Nations. 1965
Specifications for the identity and purity of food additives and their toxicological evaluation: food colors and some antimicrobials and antioxidants
World Health Organization Tech. Rep. Ser. No. 309 p. 15-16, 25
- * 516 World Health Organization of the United Nations. 1968
Specifications for the identity and purity of food additives and their toxicological evaluation: some flavouring substances and non-nutritive sweetening agents
World Health Organization Tech. Rep. Ser. No. 383 p. 12, 18
- 517 Yahiro, Arthur T. 1959
In vitro studies of formic and acetic acid metabolism by bovine rumen microorganisms
Diss. Absts. 19(12):3090
- 518 Yang, Shang-Pa. 1969
Simple method for assay of formic acid-C-14
Anal. Biochem. 32(3):519-521
- 519 York, L.R. 1971
Microbial counts and organic acid quantitation as quality indices of egg products
Diss. Abstr. Int. Sect. B. Sci. Eng. 32(6):3427, Order No. 71-31338
- 520 Yosii, Syunitiro. 1937
Beitrage zur kenntnis der fettsaureoxydation im tierkorper
J. Biochem. 26:197-224
- 521 Yuck, M.H., and F.M. Strong. 1960
Some volatile constituents of cooked beef
J. Agric. Food Chem. 8:491-494
- 522 Zakowska, Z. 1970
Estimation of volatile acids in beet molasses
Rocz. Technol. Chem. Zyrn. 19:37-45
- 523 Zhonzhorova, O.D. 1969
Studies on some volatile substances and lipid fractions formed during ripening of the Bulgarian White pickled and Kachkaval cheese
Milchwissenschaft 24(12):734-736
- 524 Zimmermann, Edmund. 1935
Die baktericide wirkung organischer sauren gegenuber bangbacillen
Zbl. Bakt. Parasitenk. Infektionskr. Abt. I. 134:215-221